

Asymptotic Theory of Bayes Factor in Stochastic Differential Equations: Part I

Trisha Maitra and Sourabh Bhattacharya*

Abstract

Research on asymptotic model selection in the context of stochastic differential equations (*SDE*'s) is almost non-existent in the literature. In particular, when a collection of *SDE*'s is considered, the problem of asymptotic model selection has not been hitherto investigated. Indeed, even though the diffusion coefficients may be considered known, questions on appropriate choice of the drift functions constitute a non-trivial model selection problem.

In this article, we develop the asymptotic theory for comparisons between collections of *SDE*'s with respect to the choice of drift functions using Bayes factors when the number of equations (individuals) in the collection of *SDE*'s tend to infinity while the time domains remain bounded for each equation. Our asymptotic theory covers situations when the observed processes associated with the *SDE*'s are independently and identically distributed (*iid*), as well as when they are independently but not identically distributed (*non-iid*). In particular, we allow incorporation of available time-dependent covariate information into each *SDE* through a multiplicative factor of the drift function; we also permit different initial values and domains of observations for the *SDE*'s.

Our model selection problem thus encompasses selection of a set of appropriate time-dependent covariates from a set of available time-dependent covariates, besides selection of the part of the drift function free of covariates.

For both *iid* and *non-iid* set-ups we establish almost sure exponential convergence of the Bayes factor.

Furthermore, we demonstrate with simulation studies that even in non-asymptotic scenarios Bayes factor successfully captures the right set of covariates.

Keywords: *Bayes factor consistency; Kullback-Leibler divergence; Martingale; Stochastic differential equations; Time-dependent covariates; Variable selection.*

1 Introduction

In statistical applications where “within” subject variability is caused by some random component varying continuously in time, stochastic differential equations (*SDE*'s) have important roles to play for modeling the temporal component of each individual. The inferential abilities of the *SDE*'s can be enhanced by incorporating covariate information available for the subjects. In these time-dependent situations it is only natural that the available covariates are also continuously varying with time. Examples of statistical applications of *SDE*-based models with time-dependent covariates are Oravecz *et al.* (2011), Overgaard *et al.* (2005), Leander *et al.* (2015), the first one also considering the hierarchical Bayesian paradigm.

Unfortunately, asymptotic inference in systems of *SDE* based models consisting of time-varying covariates seem to be rare in the statistical literature, in spite of their importance. So far random effects *SDE* models have been considered for asymptotic inference, without covariates. We refer to Delattre *et al.* (2013) for a brief review, who also undertake theoretical and classical asymptotic investigation of a class of random effects models based on *SDE*'s. Specifically, they model the i -th individual by

$$dX_i(t) = b(X_i(t), \phi_i)dt + \sigma(X_i(t))dW_i(t), \quad (1.1)$$

where, for $i = 1, \dots, n$, $X_i(0) = x^i$ is the initial value of the stochastic process $X_i(t)$, which is assumed to be continuously observed on the time interval $[0, T_i]$; $T_i > 0$ assumed to be known. The function $b(x, \varphi)$, which is the drift function, is a known, real-valued function on $\mathbb{R} \times \mathbb{R}^d$ ($\mathbb{R}$ is the real line and d is the dimension), and the function $\sigma : \mathbb{R} \mapsto \mathbb{R}$ is the known diffusion coefficient. The

*Trisha Maitra is a PhD student and Sourabh Bhattacharya is an Associate Professor in Interdisciplinary Statistical Research Unit, Indian Statistical Institute, 203, B. T. Road, Kolkata 700108. Corresponding e-mail: sourabh@isical.ac.in.

SDE's given by (1.1) are driven by independent standard Wiener processes $\{W_i(\cdot); i = 1, \dots, n\}$, and $\{\phi_i; i = 1, \dots, n\}$, which are to be interpreted as the random effect parameters associated with the n individuals, which are assumed by Delattre *et al.* (2013) to be independent of the Brownian motions and independently and identically distributed (*iid*) random variables with some common distribution. For the sake of convenience Delattre *et al.* (2013) (see also Maitra and Bhattacharya (2016c) and Maitra and Bhattacharya (2015)) assume $b(x, \phi_i) = \phi_i b(x)$. Thus, the random effect is a multiplicative factor of the drift function. In this article, we generalize the multiplicative factor to include time-dependent covariates.

In the case of *SDE*-based models, proper specification of the drift function and the associated prior distributions demand serious attention, and this falls within the purview of model selection. Moreover, when (time-varying) covariate information are available, there arises the problem of variable selection, that is, the most appropriate subset from the set of many available covariates needs to be chosen. As is well-known (see, for example, Kass and Raftery (1995)), the Bayes factor (Jeffreys (1961)) is a strong candidate for dealing with complex model selection problems. Hence, it is natural to consider this criterion for model selection in *SDE* set-ups. However, dealing with Bayes factors directly in *SDE* set-ups is usually infeasible due to unavailability of closed form expressions, and hence various numerical approximations based on Markov chain Monte Carlo, as well as related criteria such as Akaike Information Criterion (Akaike (1973)) and Bayes Information Criterion (Schwarz (1978)), are generally employed (see, for example, Fuchs (2013), Iacus (2008)). But quite importantly, although Bayes factor and its variations find use in general *SDE* models, in our knowledge covariate selection in *SDE* set-ups has not been addressed so far.

Moreover, asymptotic theory of Bayes factors in *SDE* contexts, with or without covariates, is still lacking (but see Sivaganesan and Lingham (2002) who asymptotically compare three specific diffusion models in single equation set-ups using intrinsic and fractional Bayes factors). In this paper, our goal is to develop an asymptotic theory of Bayes factors for comparing different sets of *SDE* models. Our asymptotic theory simultaneously involves time-dependent covariate selection associated with a multiplicative part of the drift function, in addition to selection of the part of the drift function free of covariates. The asymptotic framework of this paper assumes that the number of individuals tends to infinity, while their domains of observations remain bounded.

It is important to clarify that the diffusion coefficient is not associated with the question of model selection. Indeed, it is already known from Roberts and Stramer (2001) that when the associated continuous process is completely observed, the diffusion coefficient of the relevant *SDE* can be calculated directly. Moreover, two diffusion processes with different diffusion coefficients are orthogonal. Consequently, we assume throughout that the diffusion coefficient of the *SDE*'s is known.

We first develop the model selection theory using Bayes factor in general *SDE* based *iid* set-up; note that the *iid* set-up ensues when there is no covariate associated with the model and when the initial values and the domains of observations are the same for every individual. The model selection problem in *iid* cases is essentially associated with the choice of the drift functions with no involvement of covariate selection. We then extend our theory to the non-*iid* set-up, consisting of time-varying covariates and different initial values and domains of observations. Here model selection involves not only selection of the part of the drift functions free of the covariates, but also the subset of important covariates from a set of available covariates.

Specifically, we prove almost sure exponential convergence of the relevant Bayes factors in our set-ups. Assuming the *iid* set-up we develop our asymptotic theory based on a general result already existing in the literature. However, for the non-*iid* situation we first develop a general theorem which may perhaps be of independent interest, and prove almost sure exponential convergence of the Bayes factor in our non-*iid* *SDE* set-up as a special case of our theorem.

It is important to note that (which we also clarify subsequently in Section 2.6), that in the asymptotic framework of this paper, where the domains of observations remain bounded for the individuals, incorporation of random effects does not make sense from the asymptotic perspective. For this reason we include random effects in our paper Maitra and Bhattacharya (2016b), where we assume that even

the domains of observations are allowed to increase indefinitely.

The rest of our article is structured as follows. In Section 2 we formalize the problem of model selection in our aforementioned asymptotic framework. We then present the necessary assumptions and results in Section 3. In Section 4 we investigate convergence of the Bayes factor when the *SDE* models being compared form an *iid* system of equations. In Section 5 we develop a general asymptotic theory of Bayes factors in the non-*iid* situation, and then in Section 6 we investigate exponential convergence of the Bayes factor when the system of *SDE*'s are non-*iid*. In Section 7 we demonstrate with simulation studies that Bayes factor yields the correct covariate combination for our *SDE* models even in non-asymptotic cases. We provide a brief summary of this article and make concluding remarks in Section 8.

The proofs of our lemmas and theorems are provided in the supplementary document Maitra and Bhattacharya (2016d), whose sections will be referred to in this article by the prefix “S-”.

2 Formalization of the model selection problem in the *SDE* set-up

Our assumptions (H2') in Section 3 ensure that our considered systems are well defined and we are able to compute the exact likelihood. We consider the filtration $(\mathcal{F}_t^W, t \geq 0)$, where $\mathcal{F}_t^W = \sigma(W_i(s), s \leq t)$. Each process W_i is a $(\mathcal{F}_t^W, t \geq 0)$ -adapted Brownian motion.

In connection with model selection we must analyze the same data set with respect to two different models. So, although the distribution of the underlying stochastic process under the two models are different, to avoid notational complexity we denote the process by $X_i(t)$ under both the models, keeping in mind that the distinction becomes clear from the context and also by the model-specific parameters.

2.1 The structure of the *SDE* models to be compared

Now, let us consider the following two systems of *SDE* models for $i = 1, 2, \dots, n$:

$$dX_i(t) = \phi_{i,\xi_0}(t)b_{\beta_0}(X_i(t))dt + \sigma(X_i(t))dW_i(t) \quad (2.1)$$

and

$$dX_i(t) = \phi_{i,\xi_1}(t)b_{\beta_1}(X_i(t))dt + \sigma(X_i(t))dW_i(t) \quad (2.2)$$

where, $X_i(0) = x^i$ is the initial value of the stochastic process $X_i(t)$, which is assumed to be continuously observed on the time interval $[0, T_i]$; $T_i > 0$ for all i and assumed to be known. We assume that (2.1) represents the true model and (2.2) is any other model. In the above equations, for $j = 0, 1$, ξ_j and β_j denote the sets of parameters associated with the true model and the other model.

2.2 Incorporation of time-varying covariates

We model $\phi_{i,\xi_j}(t)$ for $j = 0, 1$, as

$$\phi_{i,\xi_j}(t) = \phi_{i,\xi_j}(z_i(t)) = \xi_{0j} + \xi_{1j}g_1(z_{i1}(t)) + \xi_{2j}g_2(z_{i2}(t)) + \dots + \xi_{pj}g_p(z_{ip}(t)), \quad (2.3)$$

where $\xi_j = (\xi_{0j}, \xi_{1j}, \dots, \xi_{pj})$ is a set of real constants for $j = 0, 1$, and $z_i(t) = (z_{i1}(t), z_{i2}(t), \dots, z_{ip}(t))$ is the set of available covariate information corresponding to the i -th individual, depending upon time t . We assume $z_i(t)$ is continuous in t , $z_{il}(t) \in \mathbf{Z}_l$ where $\mathbf{Z}_l$ is compact and $g_l : \mathbf{Z}_l \rightarrow \mathbb{R}$ is continuous, for $l = 1, \dots, p$. We let $\mathbf{Z} = \mathbf{Z}_1 \times \dots \times \mathbf{Z}_p$, and $\mathfrak{Z} = \{z(t) \in \mathbf{Z} : t \in [0, \infty) \text{ such that } z(t) \text{ is continuous in } t\}$. Hence, $z_i \in \mathfrak{Z}$ for all i . The functions b_{β_j} are multiplicative parts of the drift functions free of the covariates.

2.3 Model selection with respect to the drift function and the covariates

We accommodate the possibility that the dimensions of β_0, β_1 , associated with the drift functions, may be different. In reality, b_{β_0} may be piecewise linear or convex combinations of linear functions, where the number of linear functions involved (and hence, the number of associated intercept and slope parameters) may be unknown. That is, not only the values of the components of the parameter β_0 , but also the number of the components of β_0 may be unknown in reality. In general, b_{β_0} may be any function, linear or non-linear, satisfying some desirable conditions. Linearity assumptions may be convenient, but need not necessarily be unquestionable. In other words, modeling b_{β_0} in the *SDE* context is a challenging exercise, and hence the issue of model selection in this context must play an important role in the *SDE* set-up.

We also accommodate the possibility that ξ_0 and ξ_1 , associated with ϕ_{i,ξ_0} and ϕ_{i,ξ_1} , may be coefficients associated with different subsets of the available set of p covariates. This has important implication from the viewpoint of variable selection. Indeed, in a set of p time-dependent covariates, all the covariates are unlikely to be significant, particularly if p is large. Thus, some (perhaps, many) of the coefficients ξ_{i0} associated with the true model must be zero. This means that only a specific subset of the p covariates is associated with the true model. If a different set of covariates, associated with ξ_1 , is selected for actually modeling the data, then the Bayes factor is expected to favour the true set of covariates associated with ξ_0 .

If two different models are compared by the Bayes factor, none of which may be the true model, then the Bayes factor is expected to favour that model which is closest to the true model in terms of the Kullback-Leibler divergence.

2.4 Form of the Bayes factor

For $j = 0, 1$, letting $\theta_j = (\beta_j, \xi_j)$, we first define the following quantities:

$$U_{i,\theta_j} = \int_0^{T_i} \frac{\phi_{i,\xi_j}(s)b_{\beta_j}(X_i(s))}{\sigma^2(X_i(s))} dX_i(s), \quad V_{i,\theta_j} = \int_0^{T_i} \frac{\phi_{i,\xi_j}^2(s)b_{\beta_j}^2(X_i(s))}{\sigma^2(X_i(s))} ds \quad (2.4)$$

for $j = 0, 1$ and $i = 1, \dots, n$.

Let $\mathcal{C}_{T_i}$ denote the space of real continuous functions $(x(t), t \in [0, T_i])$ defined on $[0, T_i]$, endowed with the σ -field $\mathcal{C}_{T_i}$ associated with the topology of uniform convergence on $[0, T_i]$. We consider the distribution $P_j^{x_i, T_i, z_i}$ on $(\mathcal{C}_{T_i}, \mathcal{C}_{T_i})$ of $(X_i(t), t \in [0, T_i])$ given by (2.1) and (2.2) for $j = 0, 1$. We choose the dominating measure P_i as the distribution of (2.1) and (2.2) with null drift. So, for $j = 0, 1$,

$$\frac{dP_j^{x_i, T_i, z_i}}{dP_i} = f_{i,\theta_j}(X_i) = \exp \left(U_{i,\theta_j} - \frac{V_{i,\theta_j}}{2} \right), \quad (2.5)$$

where $f_{i,\theta_0}(X_i)$ denotes the true density and $f_{i,\theta_1}(X_i)$ stands for the other density associated with the modeled *SDE*.

Let $\Theta = \mathfrak{B} \times \Gamma$ be the parameter space on which a prior probability measure of θ_1 , which we denote by $\pi(\theta_1)$, is proposed. In the set-up where $n \rightarrow \infty$ and T_i are given, we are interested in asymptotic properties of the Bayes factor, given by, $I_0 \equiv 1$ and for $n \geq 1$,

$$I_n = \int_{\Theta} R_n(\theta_1) \pi(d\theta_1), \quad (2.6)$$

as $n \rightarrow \infty$, where

$$R_n(\theta_1) = \prod_{i=1}^n \frac{f_{i,\theta_1}(X_i)}{f_{i,\theta_0}(X_i)}.$$

2.5 The *iid* and the non-*iid* set-ups

Note that, for *iid* set-up $\theta_j = (\beta_j, \xi_{0j})$, along with $x^i = x$ and $T_i = T$ for all i . Since, for the *iid* set-up $\xi_j = \xi_{0j}$, so, in this case $\Gamma = \mathbb{R}$. Thus, here the problem of model selection reduces to comparing $\xi_{00}b_{\beta_0}$ with $\xi_{01}b_{\beta_1}$ using Bayes factor.

In the non-*iid* set-up we relax the assumptions $\xi_{1j} = \xi_{2j} = \dots = \xi_{pj} = 0$ and $x^i = x$, $T_i = T$ for each i . Hence, in this case, the model selection problem involves variable selection as well as comparison between different drift functions.

2.6 No random effects when T_i are given

It is important to perceive that when the T_i are fixed constants, it is not possible to allow random effects into the model and still achieve consistency of the Bayes factor. This is because in that case the *SDE* set-up would simply reduce to n independent models, each with independent sets of parameters, leaving no scope for asymptotics since T_i are held constants. In Maitra and Bhattacharya (2016b) we consider random effects when $T_i \rightarrow \infty$ along with $n \rightarrow \infty$.

2.7 A key relation between U_{i,θ_j} and V_{i,θ_j} in the context of model selection using Bayes factors

An useful relation between U_{i,θ_j} and V_{i,θ_j} which we will often make use of in this paper is as follows.

$$\begin{aligned} U_{i,\theta_j} &= \int_0^{T_i} \frac{\phi_{i,\xi_j}(s)b_{\beta_j}(X_i(s))}{\sigma^2(X_i(s))} dX_i(s) \\ &= \int_0^{T_i} \frac{\phi_{i,\xi_j}(s)b_{\beta_j}(X_i(s))}{\sigma^2(X_i(s))} [\phi_{i,\xi_0}(s)b_{\beta_0}(X_i(s)) ds + \sigma(X_i(s)) dW_i(s)] \\ &= \int_0^{T_i} \frac{\phi_{i,\xi_j}(s)\phi_{i,\xi_0}(s)b_{\beta_j}(X_i(s))b_{\beta_0}(X_i(s))}{\sigma^2(X_i(s))} ds + \int_0^{T_i} \frac{\phi_{i,\xi_j}(s)b_{\beta_j}(X_i(s))}{\sigma(X_i(s))} dW_i(s) \\ &= V_{i,\theta_0,\theta_j} + \int_0^{T_i} \frac{\phi_{i,\xi_j}(s)b_{\beta_j}(X_i(s))}{\sigma(X_i(s))} dW_i(s), \end{aligned} \quad (2.7)$$

with

$$V_{i,\theta_0,\theta_j} = \int_0^{T_i} \frac{\phi_{i,\xi_j}(s)\phi_{i,\xi_0}(s)b_{\beta_j}(X_i(s))b_{\beta_0}(X_i(s))}{\sigma^2(X_i(s))} ds. \quad (2.8)$$

Note that $V_{i,\theta_0} = V_{i,\theta_0,\theta_0}$ and $V_{i,\theta_1} = V_{i,\theta_1,\theta_1}$. Also note that, for $j = 0, 1$, for each i ,

$$E_{\theta_0} \left[\int_0^{T_i} \frac{\phi_{i,\xi_j}(s)b_{\beta_j}(X_i(s))}{\sigma(X_i(s))} dW_i(s) \right] = 0, \quad (2.9)$$

so that $E_{\theta_0}(U_{i,\theta_j}) = E_{\theta_0}(V_{i,\theta_0,\theta_j})$.

3 Requisite assumptions and results for the asymptotic theory of Bayes factor when $n \rightarrow \infty$ but T_i are constants for every i

We assume the following conditions:

(H1') The parameter space $\Theta = \mathfrak{B} \times \Gamma$ such that $\mathfrak{B}$ and Γ are compact.

(H2') For $j = 0, 1$, $b_{\beta_j}(\cdot)$ and $\sigma(\cdot)$ are C^1 on $\mathbb{R}$ and satisfy $b_{\beta_j}^2(x) \leq K_1(1 + x^2 + \|\beta_j\|^2)$ and $\sigma^2(x) \leq K_2(1 + x^2)$ for all $x \in \mathbb{R}$, for some $K_1, K_2 > 0$. Now, due to (H1') the latter boils down to assuming $b_{\beta_j}^2(x) \leq K(1 + x^2)$ and $\sigma^2(x) \leq K(1 + x^2)$ for all $x \in \mathbb{R}$, for some $K > 0$.

Because of (H2') it follows from Theorem 4.4 of Mao (2011), page 61, that for all $T_i > 0$, and any $k \geq 2$,

$$E \left(\sup_{s \in [0, T_i]} |X_i(s)|^k \right) \leq \left(1 + 3^{k-1} E |X_i(0)|^k \right) \exp \left(\tilde{\vartheta} T_i \right), \quad (3.1)$$

where

$$\tilde{\vartheta} = \frac{1}{6} (18K)^{\frac{k}{2}} T_i^{\frac{k-2}{2}} \left[T_i^{\frac{k}{2}} + \left(\frac{k^3}{2(k-1)} \right)^{\frac{k}{2}} \right].$$

We further assume:

(H3') For every x , $b_{\beta_j}(x)$ is continuous in β_j , for $j = 0, 1$.

(H4') For $j = 0, 1$,

$$\frac{b_{\beta_j}^2(x)}{\sigma^2(x)} \leq K_{\beta_j} (1 + x^2 + \|\beta_j\|^2), \quad (3.2)$$

where K_{β_j} is continuous in β_j .

(H5') (i) Let $\mathcal{Z} = \mathbf{Z}_1 \times \mathbf{Z}_2 \times \cdots \times \mathbf{Z}_p$ be the space of covariates where $\mathbf{Z}_l$ is compact for $l = 1, \dots, p$ and $\mathbf{z}_i(t) = (z_{i1}(t), z_{i2}(t), \dots, z_{ip}(t)) \in \mathcal{Z}$ for every $i = 1, \dots, n$ and $t \in [0, T_i]$. Moreover, $\mathbf{z}_i(t)$ are continuous in t , so that $\mathbf{z}_i \in \mathfrak{Z}$ for every i .

(ii) For $j = 0, 1$, the vector of covariates $\mathbf{z}_i(t)$ is related to the i -th SDE of the j -th model via

$$\phi_{i, \xi_j}(t) = \phi_{\xi_j}(\mathbf{z}_i(t)) = \xi_{0j} + \sum_{l=1}^p \xi_{lj} g_l(\mathbf{z}_i(t)),$$

where, for $l = 1, \dots, p$, $g_l : \mathbf{Z}_l \rightarrow \mathbb{R}$ is continuous. Notationally, for a given $\mathbf{z}(t)$, we denote $\phi_{\xi_j}(t) = \phi_{\xi_j}(\mathbf{z}(t)) = \xi_{0j} + \sum_{l=1}^p \xi_{lj} g_l(\mathbf{z}(t))$.

(iii) For $l = 1, \dots, p$, and for $t \in [0, T_i]$,

$$\frac{1}{n} \sum_{i=1}^n g_l(z_{il}(t)) \rightarrow c_l(t); \quad (3.3)$$

and, for $l, m = 1, \dots, p$; $t \in [0, T_i]$,

$$\frac{1}{n} \sum_{i=1}^n g_l(z_{il}(t)) g_m(z_{im}(t)) \rightarrow c_l(t) c_m(t), \quad (3.4)$$

as $n \rightarrow \infty$, where $c_l(t)$ are real constants.

Note that, given l and t , had $z_{il}(t)$ been random and *iid* with respect to i , then (3.3) would hold almost surely by the strong law of large numbers. Additionally, if $z_{il}(t)$ and $z_{im}(t)$ were independent, then (3.4) would hold almost surely as well. Hence, in this paper, one may assume that for $i = 1, \dots, n$, and $l = 1, \dots, p$, the covariates z_{il} are observed realizations of stochastic processes that are *iid* for $i = 1, \dots, n$, for all $l = 1, \dots, p$, and that for $l \neq m$, the processes generating z_{il} and z_{im} are independent. Thus, in essence, we assume here that for $l \neq m$, $g_l(z_{il}(t))$ and $g_m(z_{im}(t))$ are uncorrelated.

We then have the following lemma, which will be useful for proving our main results.

Lemma 1 Assume (H1') – (H4'). Then for all $\theta_1 \in \mathfrak{B} \times \Gamma$, for $k \geq 1$,

$$E_{\theta_0} [U_{i, \theta_j}]^k < \infty; j = 0, 1, \quad (3.5)$$

$$E_{\theta_0} [V_{i, \theta_1}]^k < \infty, \quad (3.6)$$

$$E_{\theta_0} [V_{i, \theta_0, \theta_j}]^k < \infty; j = 0, 1. \quad (3.7)$$

Moreover, for $j = 1$, the above expectations are continuous in θ_1 .

4 Convergence of Bayes factor in the *SDE* based *iid* set-up

We first consider the *iid* set-up; in other words, we assume that $x^i = x$, $T_i = T$ for $i = 1, \dots, n$, and $j = 0, 1$. In this case $\theta_j = (\beta_j, \xi_{0j})$ for $j = 0, 1$. We shall relax these assumptions subsequently when we take up the non-*iid* (that is, independent, but non-identical) case.

4.1 A general result on consistency of Bayes factor in the *iid* set-up

To investigate consistency of the Bayes factor, we resort to a general result in the *iid* set-up developed by Walker (2004) (see also Walker *et al.* (2004)). To state the result we first define some relevant notation which apply to both parametric and nonparametric problems. For any x in the appropriate domain, let

$$\hat{f}_n(x) = \int f(x) \pi_n(df)$$

be the posterior predictive density, where π_n stands for the posterior of f , given by

$$\pi_n(A) = \frac{\int_A \prod_{i=1}^n f(X_i) \pi(df)}{\int \prod_{i=1}^n f(X_i) \pi(df)}$$

and let

$$\hat{f}_{nA}(x) = \int f(x) \pi_{nA}(df)$$

be the posterior predictive density restricted to the set A , that is, for the prior probability $\pi(A) > 0$,

$$\pi_{nA}(df) = \frac{I_A(f) \pi_n(df)}{\int_A \pi_n(df)},$$

where I_A denotes the indicator function of the set A .

Clearly, the above set-up is in accordance with the *iid* situation. The following theorem of Walker (2004) is appropriate for our *iid* set-up.

Theorem 2 (Walker (2004)) Assume that

$$\pi(f : \mathcal{K}(f, f_0) < c_1) > 0, \quad (4.1)$$

only for, and for all $c_1 > \delta$, for some $\delta \geq 0$, and that for all $\epsilon > 0$,

$$\liminf_n \mathcal{K}(f_0, \hat{f}_{nA(\epsilon)}) \geq \epsilon, \quad (4.2)$$

when $A(\epsilon) = \{f : \mathcal{K}(f_0, f) > \epsilon\}$. Property (4.1) is the Kullback-Leibler property and (4.2) has been referred to as the Q^* property by Walker (2004). Assume further that

$$\sup_n \text{Var} \left(\frac{I_{n+1}}{I_n} \right) < \infty. \quad (4.3)$$

Then,

$$n^{-1} \log(I_n) \rightarrow -\delta, \quad (4.4)$$

almost surely.

The following corollary provides the result on asymptotic comparison between two models using Bayes factors, in the *iid* case.

Corollary 3 (Walker (2004)) Let $R_n(f) = \prod_{i=1}^n \frac{f(X_i)}{f_0(X_i)}$. For $j = 1, 2$, let

$$I_{jn} = \int R_n(f) \pi_j(df),$$

where π_1 and π_2 are two different priors on f . Let $B_n = I_{1n}/I_{2n}$ denote the Bayes factor for comparing the two models associated with π_1 and π_2 . If π_1 and π_2 have the Kullback-Leibler property (4.1) with $\delta = \delta_1$ and $\delta = \delta_2$ respectively, satisfy the Q^* property (4.2), and (4.3) with $I_n = I_{jn}$, for $j = 1, 2$, then

$$n^{-1} \log B_n \rightarrow \delta_2 - \delta_1,$$

almost surely.

Remark 4 In Walker (2004) the densities are assumed to be dominated by the Lebesgue measure. However, this is not necessary. The results remain true if the densities are with respect to any valid measure; see, for example, Barron et al. (1999) for related concepts and results (Lemma 4 in particular) with respect to general measures. As such, in our SDE-based situation, although the densities are not dominated by the Lebesgue measure (see (2.5)), all our results still remain valid.

4.2 Verification of Theorem 2 in iid SDE set-up

In our parametric case, $f_0 \equiv f_{\theta_0}$ and $f \equiv f_{\theta_1}$. In this iid set-up, as mentioned earlier $\xi_j = \xi_{0j}$ for $j = 0, 1$, so that $\phi_{\xi_j} \equiv \xi_{0j}$. For our convenience, we let, for $j = 0, 1$ and $i = 1, \dots, n$,

$$\tilde{U}_{i,\beta_j} = \int_0^{T_i} \frac{b_{\beta_j}(X_i(s))}{\sigma^2(X_i(s))} dX_i(s), \quad \tilde{V}_{i,\beta_0,\beta_j} = \int_0^{T_i} \frac{b_{\beta_0}(X_i(s))b_{\beta_j}(X_i(s))}{\sigma^2(X_i(s))} ds. \quad (4.5)$$

Note that, for $i = 1, \dots, n$, $\tilde{V}_{i,\beta_0} = \tilde{V}_{i,\beta_0,\beta_0}$ and $\tilde{V}_{i,\beta_1} = \tilde{V}_{i,\beta_1,\beta_1}$. The Kullback-Leibler divergence measure between f_0 and f in this set-up is given, with $i = 1$, by

$$\mathcal{K}(f_{\theta_0}, f_{\theta_1}) = \frac{\xi_{00}^2}{2} E_{\theta_0}(\tilde{V}_{1,\beta_0}) - \xi_{00}\xi_{01} E_{\theta_0}(\tilde{V}_{1,\beta_0,\beta_1}) + \frac{\xi_{01}^2}{2} E_{\theta_0}(\tilde{V}_{1,\beta_1}), \quad (4.6)$$

where $E_{\theta_0} \equiv E_{f_{\theta_0}}$. The result easily follows from (2.7) and (2.9). Now let

$$\begin{aligned} \delta &= \min_{\Theta} \mathcal{K}(f_{\theta_0}, f_{\theta_1}) \\ &= \min_{\Theta} \left\{ \frac{\xi_{00}^2}{2} E_{\theta_0}(\tilde{V}_{1,\beta_0}) - \xi_{00}\xi_{01} E_{\theta_0}(\tilde{V}_{1,\beta_0,\beta_1}) + \frac{\xi_{01}^2}{2} E_{\theta_0}(\tilde{V}_{1,\beta_1}) \right\}. \end{aligned} \quad (4.7)$$

Since $E_{\theta_0}(\tilde{V}_{1,\beta_1})$ and $E_{\theta_0}(\tilde{V}_{1,\beta_0,\beta_1})$ are continuous in β_1 , compactness of Θ guarantees that $0 \leq \delta < \infty$.

4.2.1 Verification of (4.1)

To see that (4.1) holds in our case for any prior dominated by Lebesgue measure, first let us define

$$\mathcal{K}^*(f_{\bar{\theta}}, f_{\theta_1}) = \mathcal{K}(f_{\theta_0}, f_{\theta_1}) - \mathcal{K}(f_{\theta_0}, f_{\bar{\theta}}), \quad (4.8)$$

where $f_{\bar{\theta}} = \arg \min_{\Theta} \mathcal{K}(f_{\theta_0}, f_{\theta_1})$. Now, let us choose any prior π such that $\frac{d\pi}{d\nu} = \varrho$ where ϱ is a continuous positive density with respect to Lebesgue measure, where, by “positive” density, we mean a density excluding any interval of null measure. For any $c_1 > 0$, we then need to show that

$$\pi(\theta_1 \in \Theta : \delta \leq \mathcal{K}(f_{\theta_0}, f_{\theta_1}) < \delta + c_1) > 0,$$

for any prior π dominated by Lebesgue measure. This is equivalent to showing

$$\pi(\boldsymbol{\theta}_1 \in \boldsymbol{\Theta} : 0 \leq \mathcal{K}^*(f_{\bar{\boldsymbol{\theta}}}, f_{\boldsymbol{\theta}_1}) < c_1) > 0,$$

for any prior π dominated by Lebesgue measure.

Since $\mathcal{K}(f_{\boldsymbol{\theta}_0}, f_{\boldsymbol{\theta}_1})$ is continuous in $\boldsymbol{\theta}_1$, so is $\mathcal{K}^*(f_{\bar{\boldsymbol{\theta}}}, f_{\boldsymbol{\theta}_1})$. Compactness of $\boldsymbol{\Theta}$ ensures uniform continuity of $\mathcal{K}^*(f_{\bar{\boldsymbol{\theta}}}, f_{\boldsymbol{\theta}_1})$. Hence, for any $c_1 > 0$, there exists ϵ_{c_1} independent of $\boldsymbol{\theta}_1$, such that $\|\boldsymbol{\theta}_1 - \bar{\boldsymbol{\theta}}\| < \epsilon_{c_1}$ implies $\mathcal{K}^*(f_{\bar{\boldsymbol{\theta}}}, f_{\boldsymbol{\theta}_1}) < c_1$. Then,

$$\begin{aligned} \pi(\boldsymbol{\theta}_1 \in \boldsymbol{\Theta} : 0 \leq \mathcal{K}^*(f_{\bar{\boldsymbol{\theta}}}, f_{\boldsymbol{\theta}_1}) < c_1) &\geq \pi(\boldsymbol{\theta}_1 \in \boldsymbol{\Theta} : \|\boldsymbol{\theta}_1 - \bar{\boldsymbol{\theta}}\| < \epsilon_{c_1}) \\ &\geq \left[\inf_{\{\boldsymbol{\theta}_1 \in \boldsymbol{\Theta} : \|\boldsymbol{\theta}_1 - \bar{\boldsymbol{\theta}}\| < \epsilon_{c_1}\}} \varrho(\boldsymbol{\theta}_1) \right] \times \nu(\{\boldsymbol{\theta}_1 \in \boldsymbol{\Theta} : \|\boldsymbol{\theta}_1 - \bar{\boldsymbol{\theta}}\| < \epsilon_{c_1}\}) > 0, \end{aligned} \quad (4.9)$$

where ν stands for Lebesgue measure. In other words, (4.1) holds in our case.

4.2.2 Verification of (4.2)

To see that (4.2) also holds in our *SDE* set-up, first note that in our case

$$\hat{f}_{nA(\epsilon)}(x) = \frac{\int_{A(\epsilon)} f_{\boldsymbol{\theta}_1}(x) \pi_n(d\boldsymbol{\theta}_1)}{\int_{A(\epsilon)} \pi_n(d\boldsymbol{\theta}_1)}, \quad (4.10)$$

with

$$\begin{aligned} A(\epsilon) &= \{\boldsymbol{\theta}_1 \in \boldsymbol{\Theta} : \mathcal{K}(f_{\boldsymbol{\theta}_0}, f_{\boldsymbol{\theta}_1}) \geq \epsilon\} \\ &= \left\{ \boldsymbol{\theta}_1 \in \boldsymbol{\Theta} : \frac{\xi_{00}^2}{2} E_{\boldsymbol{\theta}_0}(\tilde{V}_{1,\beta_0}) - \xi_{00}\xi_{01} E_{\boldsymbol{\theta}_0}(\tilde{V}_{1,\beta_0,\beta_1}) + \frac{\xi_{01}^2}{2} E_{\boldsymbol{\theta}_0}(\tilde{V}_{1,\beta_1}) \geq \epsilon \right\} \end{aligned} \quad (4.11)$$

for any $\epsilon > 0$. Note that, here we have replaced $K(f_{\boldsymbol{\theta}_0}, f_{\boldsymbol{\theta}_1}) > \epsilon$ with $K(f_{\boldsymbol{\theta}_0}, f_{\boldsymbol{\theta}_1}) \geq \epsilon$ in the definition of $A(\epsilon)$ because of continuity of the posterior of $\boldsymbol{\theta}_1$. Note that

$$\hat{f}_{nA(\epsilon)}(X) \leq \sup_{\boldsymbol{\theta}_1 \in A(\epsilon)} f_{\boldsymbol{\theta}_1}(X) = f_{\hat{\boldsymbol{\theta}}_1(X)}(X), \quad (4.12)$$

where $\hat{\boldsymbol{\theta}}_1(X)$, which depends upon X , is the maximizer lying in the compact set $A(\epsilon)$. Now note that

$$\begin{aligned} \mathcal{K}(f_{\boldsymbol{\theta}_0}, \hat{f}_{nA(\epsilon)}) &= E_{\boldsymbol{\theta}_0} [\log f_{\boldsymbol{\theta}_0}(X)] - E_{\boldsymbol{\theta}_0} [\log \hat{f}_{nA(\epsilon)}(X)] \\ &\geq E_{\boldsymbol{\theta}_0} [\log f_{\boldsymbol{\theta}_0}(X)] - E_{\boldsymbol{\theta}_0} [\log f_{\hat{\boldsymbol{\theta}}_1(X)}(X)] \\ &= E_{\boldsymbol{\theta}_0} \left(\log \frac{f_{\boldsymbol{\theta}_0}(X)}{f_{\hat{\boldsymbol{\theta}}_1(X)}(X)} \right) \\ &= E_{\hat{\boldsymbol{\theta}}_1(X)|\boldsymbol{\theta}_0} E_{X|\hat{\boldsymbol{\theta}}_1(X)=\boldsymbol{\theta}, \boldsymbol{\theta}_0} \left(\log \frac{f_{\boldsymbol{\theta}_0}(X)}{f_{\{\hat{\boldsymbol{\theta}}_1(X)=\boldsymbol{\theta}\}}(X)} \right) \\ &= E_{\hat{\boldsymbol{\theta}}_1(X)|\boldsymbol{\theta}_0} \mathcal{K}(f_{\boldsymbol{\theta}_0}, f_{\boldsymbol{\theta}}) \\ &\geq E_{\hat{\boldsymbol{\theta}}_1(X)|\boldsymbol{\theta}_0} \inf_{\boldsymbol{\theta} \in A(\epsilon)} \mathcal{K}(f_{\boldsymbol{\theta}_0}, f_{\boldsymbol{\theta}}) \\ &= E_{\hat{\boldsymbol{\theta}}_1(X)|\boldsymbol{\theta}_0} \mathcal{K}(f_{\boldsymbol{\theta}_0}, f_{\boldsymbol{\theta}^*}) \\ &= \mathcal{K}(f_{\boldsymbol{\theta}_0}, f_{\boldsymbol{\theta}^*}) \\ &\geq \epsilon, \end{aligned} \quad (4.13)$$

where $\vartheta^* = \arg \min_{\vartheta \in A(\epsilon)} \mathcal{K}(f_{\theta_0}, f_{\vartheta})$. Hence, (4.2) is satisfied in our *SDE* set-up.

4.2.3 Verification of (4.3)

We now prove that (4.3) also holds. It is straightforward to verify that

$$\frac{I_{n+1}}{I_n} = \frac{\hat{f}_{n+1}(X_{n+1})}{f_{\theta_0}(X_{n+1})}, \quad (4.14)$$

where

$$\hat{f}_{n+1}(\cdot) = E_{\theta_1|X_1, \dots, X_n} [f_{\theta_1}(\cdot)] \quad (4.15)$$

is the posterior predictive distribution of $f_{\theta_1}(\cdot)$, with respect to the posterior of θ_1 , given $X_1, \dots, X_n$. In (4.15), $E_{\theta_1|X_1, \dots, X_n}$ denotes expectation with respect to the posterior of θ_1 given $X_1, \dots, X_n$.

First note that, since

$$\log [f_{\theta_0}(X_{n+1})] = \xi_{00} \tilde{U}_{n+1, \beta_0} - \frac{\xi_{00}^2}{2} \tilde{V}_{n+1, \beta_0}, \quad (4.16)$$

it follows from Lemma 1 that the moments of all orders of $\log [f_{\theta_0}(X_{n+1})]$ exist and are finite. Also, since X_i are *iid*, the moments are the same for every $n = 1, 2, \dots$. In other words,

$$\sup_n \text{Var} (\log f_{\theta_0}(X_{n+1})) < \infty. \quad (4.17)$$

Then observe that for any given X_{n+1} , using compactness of Θ and continuity of $f_{\theta_1}(X_{n+1})$ with respect to θ_1 ,

$$f_{\theta_1^*(X_{n+1})}(X_{n+1}) = \inf_{\theta_1 \in \Theta} f_{\theta_1}(X_{n+1}) \leq \hat{f}_{n+1}(X_{n+1}) \leq \sup_{\theta_1 \in \Theta} f_{\theta_1}(X_{n+1}) = f_{\theta_1^{**}(X_{n+1})}(X_{n+1}),$$

where $\theta_1^*(X_{n+1}) = \arg \min_{\theta_1 \in \Theta} f_{\theta_1}(X_{n+1})$ and $\theta_1^{**}(X_{n+1}) = \arg \max_{\theta_1 \in \Theta} f_{\theta_1}(X_{n+1})$. Clearly, $\theta_1^*(X_{n+1}), \theta_1^{**}(X_{n+1}) \in \Theta$, for any given X_{n+1} . Moreover,

$$\theta_1^*(X_{n+1}) = (\beta_1^*(X_{n+1}), \xi_{01}^*(X_{n+1})), \quad \theta_1^{**}(X_{n+1}) = (\beta_1^{**}(X_{n+1}), \xi_{01}^{**}(X_{n+1}))$$

where each component of $\theta_1^*(X_{n+1})$ and $\theta_1^{**}(X_{n+1})$ depends on X_{n+1} . Noting that $U_{n+1, \theta_1^*(X_{n+1})} = \xi_{01}^*(X_{n+1}) \tilde{U}_{n+1, \beta_1^*(X_{n+1})}$ and $V_{n+1, \theta_1^*(X_{n+1})} = \{\xi_{01}^*(X_{n+1})\}^2 \tilde{V}_{n+1, \beta_1^*(X_{n+1})}$, it follows from the above inequality that

$$\begin{aligned} & -|U_{n+1, \theta_1^*(X_{n+1})}| - \frac{V_{n+1, \theta_1^*(X_{n+1})}}{2} \\ & \leq U_{n+1, \theta_1^*(X_{n+1})} - \frac{V_{n+1, \theta_1^*(X_{n+1})}}{2} \\ & \leq \log \hat{f}_{n+1}(X_{n+1}) \\ & \leq U_{n+1, \theta_1^{**}(X_{n+1})} - \frac{V_{n+1, \theta_1^{**}(X_{n+1})}}{2} \\ & \leq |U_{n+1, \theta_1^{**}(X_{n+1})}| + \frac{V_{n+1, \theta_1^{**}(X_{n+1})}}{2}. \end{aligned}$$

Hence, $E_{\theta_0} \left(\log \hat{f}_{n+1}(X_{n+1}) \right)^2$ lies between $E_{\theta_0} \left(|U_{n+1, \theta_1^*(X_{n+1})}| + \frac{V_{n+1, \theta_1^*(X_{n+1})}}{2} \right)^2$ and

$$E_{\theta_0} \left(|U_{n+1, \theta_1^{**}(X_{n+1})}| + \frac{V_{n+1, \theta_1^{**}(X_{n+1})}}{2} \right)^2.$$

We obtain uniform lower and upper bounds of the above two expressions in the following manner. For the upper bound of the latter we first take supremum of the expectation with respect to X_{n+1} , conditional on $\theta_1^{**}(X_{n+1}) = \varsigma$, over $\varsigma \in \Theta$, and then take expectation with respect to X_{n+1} . Since $\varsigma \in \Theta$, compactness of Θ and Lemma 1 ensure that the moments of any given order of the above expression is uniformly bounded above. Analogously, we obtain a uniform lower bound replacing the supremum with infimum. In the same way we obtain uniform lower and upper bounds of the other expression. The uniform bounds on the second order moments, in turn, guarantee that

$$\sup_n \text{Var} \left(\log \hat{f}_{n+1}(X_{n+1}) \right) < \infty. \quad (4.18)$$

Combining (4.17) and (4.18) and using the Cauchy-Schwartz inequality for the covariance term associated with $\text{Var} \left(\log \hat{f}_{n+1}(X_{n+1}) - \log f_{\theta_0}(X_{n+1}) \right)$ shows that (4.3) holds in our set-up.

We formalize the above arguments in the form of a theorem in the *SDE* based *iid* set-up.

Theorem 5 *Assume the iid case of the SDE based set-up and conditions (H1') – (H4'). Then (4.4) holds.*

The following corollary in the *iid SDE* context is motivated by Corollary 3.

Corollary 6 *For $j = 1, 2$, let $R_{jn}(\theta_j) = \prod_{i=1}^n \frac{f_{\theta_j}(X_i)}{f_{\theta_0}(X_i)}$, where θ_1 and θ_2 are two different finite sets of parameters, perhaps with different dimensionalities, associated with the two models to be compared. For $j = 1, 2$, let*

$$I_{jn} = \int R_{jn}(\theta_j) \pi_j(d\theta_j),$$

where π_j is the prior on θ_j . Let $B_n = I_{1n}/I_{2n}$ as before. Assume the iid case of the *SDE* based set-up and suppose that both the models satisfy conditions (H1') – (H4') and have the Kullback-Leibler property with $\delta = \delta_1$ and $\delta = \delta_2$ respectively. Then

$$n^{-1} \log B_n \rightarrow \delta_2 - \delta_1,$$

almost surely.

5 General asymptotic theory of Bayes factor in the non-*iid* set-up

In this section, we first develop a general asymptotic theory of Bayes factors in the non-*iid* set-up, and then obtain the result for the non-*iid SDE* set-up as a special case of our general theory.

5.1 The basic set-up

We assume that for $i = 1, \dots, n$, $X_i \sim f_{0i}$, that is, the true density function corresponding to the i -th individual is f_{0i} . Considering another arbitrary density f_i for individual X_i we investigate consistency of the Bayes factor in this general non-*iid* set-up. For our purpose we introduce the following two properties:

1. Kullback-Leibler (δ) property in the non-*iid* set-up:

We denote the Kullback-Leibler divergence measure between f_{0i} and f_i by $\mathcal{K}(f_{0i}, f_i)$ and assume that the limit

$$\mathcal{K}^\infty(f_0, f) = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n E \left[\log \frac{f_{0i}(X_i)}{f_i(X_i)} \right] = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \mathcal{K}(f_{0i}, f_i) \quad (5.1)$$

exists almost surely with respect to the prior π on f . Let the prior distribution π satisfy

$$\pi \left(f : \inf_i \mathcal{K}(f_{0i}, f_i) \geq \delta \right) = 1, \quad (5.2)$$

for some $\delta \geq 0$. Then we say that π has the Kullback-Leibler (δ) property if, for any $c > 0$,

$$\pi(f : \delta \leq \mathcal{K}^\infty(f_0, f) \leq \delta + c) > 0. \quad (5.3)$$

2. Q^* property in the non-*iid* set-up:

Let us denote the posterior distribution corresponding to n observations by π_n . We denote $\pi(df_1, df_2, \dots, df_n)$ by $\pi(\tilde{df})$. For any set A ,

$$\pi_n(A) = \frac{\int_A \prod_{i=1}^n f_i(X_i) \pi(\tilde{df})}{\int \prod_{i=1}^n f_i(X_i) \pi(\tilde{df})}$$

denotes the posterior probability of A . Let

$$R_n(f_1, f_2, \dots, f_n) = \prod_{i=1}^n \frac{f_i(X_i)}{f_{0i}(X_i)}.$$

Let us define the posterior predictive density by

$$\hat{f}_n(X_n) = \int f_n(X_n) \pi_n(df_n),$$

and

$$\hat{f}_{nA}(X_n) = \int f_n(X_n) \pi_{nA}(df_n)$$

to be the posterior predictive density with posterior restricted to the set A , that is, for $\pi(A) > 0$,

$$\pi_{nA}(df_n) = \frac{\mathbf{I}_A(f_n) \pi_n(df_n)}{\int_A \pi_n(df_n)}.$$

Then we say that the prior has the *property Q^* in the non-*iid* set-up* if the following holds for any $\epsilon > 0$:

$$\liminf_n \mathcal{K}(f_{0n}, \hat{f}_{n, A_n(\epsilon)}) \geq \epsilon, \quad (5.4)$$

when

$$A_n(\epsilon) = \{f_n : \mathcal{K}(f_{0n}, f_n) \geq \epsilon\}. \quad (5.5)$$

Let $I_0 \equiv 1$ and for $n \geq 1$, let us define

$$I_n = \int R_n(f_1, f_2, \dots, f_n) \pi(\tilde{df}), \quad (5.6)$$

which is relevant for the study of the Bayes factors. Regarding convergence of I_n , we formulate the following theorem.

Theorem 7 *Assume the non-*iid* set-up and that the limit (5.1) exists almost surely with respect to the prior π . Also assume that the prior π satisfies (5.2), has the Kullback-Leibler (δ) and Q^* properties given by (5.3) and (5.4), respectively. Assume further that*

$$\sup_i E \left[\log \frac{f_{0i}(X_i)}{f_i(X_i)} \right]^2 < \infty \quad (5.7)$$

and

$$\sup_n E \left[\log \frac{I_n}{I_{n-1}} \right]^2 < \infty. \quad (5.8)$$

Then

$$n^{-1} \log I_n \rightarrow -\delta, \quad (5.9)$$

almost surely as $n \rightarrow \infty$.

Corollary 8 For $j = 1, 2$, let

$$I_{jn} = \int R_n(f_1, \dots, f_n) \pi_j(\tilde{d}f),$$

where π_1 and π_2 are two different priors on f . Let $B_n = I_{1n}/I_{2n}$ denote the Bayes factor for comparing the two models associated with π_1 and π_2 . If both the models satisfy the conditions of Theorem 7, and satisfy the Kullback-Leibler property with $\delta = \delta_1$ and $\delta = \delta_2$ respectively, then

$$n^{-1} \log B_n \rightarrow \delta_2 - \delta_1,$$

almost surely.

6 Specialization of non-*iid* asymptotic theory of Bayes factors to non-*iid* SDE set-up where T_i are constants for every i but $n \rightarrow \infty$

In this section we relax the restrictions $T_i = T$ and $x^i = x$ for $i = 1, \dots, n$. In other words, here we deal with the set-up where the processes $X_i(\cdot)$; $i = 1, \dots, n$, are independently, but not identically distributed. Following Maitra and Bhattacharya (2016c), Maitra and Bhattacharya (2015) we assume the following:

(H6') The sequences $\{T_1, T_2, \dots\}$ and $\{x^1, x^2, \dots\}$ are sequences in compact sets $\mathfrak{T}$ and $\mathfrak{X}$, respectively, so that there exist convergent subsequences with limits in $\mathfrak{T}$ and $\mathfrak{X}$. For notational convenience, we continue to denote the convergent subsequences as $\{T_1, T_2, \dots\}$ and $\{x^1, x^2, \dots\}$. Let us denote the limits by T^∞ and x^∞ , where $T^\infty \in \mathfrak{T}$ and $x^\infty \in \mathfrak{X}$.

Remark 9 Note that the choices of the convergent subsequences $\{T_1, T_2, \dots\}$ and $\{x^1, x^2, \dots\}$ are not unique. However, this non-uniqueness does not affect asymptotic selection of the correct model via Bayes factor. Indeed, as will be evident from our proof, for any choice of convergent subsequence, the Bayes factor almost surely converges exponentially to the correct quantity. The reason for this is that we actually need to deal with the infimum of the Kullback-Leibler distance over $\mathfrak{X}$ and $\mathfrak{T}$, which is of course independent of the choices of subsequences; see Section 6.1 for the details.

Following Maitra and Bhattacharya (2016c), we denote the process associated with the initial value x and time point t as $X(t, x)$, so that $X(t, x^i) = X_i(t)$, and $X_i = \{X_i(t); t \in [0, T_i]\}$.

Let $\theta_j = (\beta_j, \xi_j)$ for $j = 0, 1$ denote the set of finite number of parameters, where β_j and ξ_j have the same interpretation as in the *iid* set-up. As before, $z_i(t) = (z_{i1}(t), z_{i2}(t), \dots, z_{ip}(t))$ is the set of covariate information corresponding to i -th individual at time point t . For $x^i \in \mathfrak{X}$, $T_i \in \mathfrak{T}$, $z_i(t) \in \mathfrak{Z}$ and $\theta_j \in \Theta$, let

$$U_{x^i, T_i, z_i, \theta_j} = \int_0^{T_i} \frac{\phi_{i, \xi_j}(s) b_{\beta_j}(X_i(s, x^i))}{\sigma^2(X_i(s, x^i))} dX_i(s, x^i); \quad (6.1)$$

$$V_{x^i, T_i, z_i, \theta_0, \theta_j} = \int_0^{T_i} \frac{\phi_{i, \xi_j}(s) \phi_{i, \xi_0}(s) b_{\beta_j}(X_i(s, x^i)) b_{\beta_0}(X_i(s, x^i))}{\sigma^2(X_i(s, x^i))} ds. \quad (6.2)$$

As before, $V_{x^i, T_i, z_i, \theta_0} = V_{x^i, T_i, z_i, \theta_0, \theta_0}$ and $V_{x^i, T_i, z_i, \theta_1} = V_{x^i, T_i, z_i, \theta_1, \theta_1}$.

In this non-*iid* set-up $f_{0i} = f_{\theta_0, x^i, T_i, z_i}$ and $f_i = f_{\theta_1, x^i, T_i, z_i}$. An extension of Lemma 1 incorporating x, T and z shows that moments of U_{x, T, z, θ_j} , V_{x, T, z, θ_j} , $V_{x, T, z, \theta_0, \theta_j}$ of all orders exist, and are continuous in x, T, z, θ_1 . Formally, we have the following lemma.

Lemma 10 Assume (H1') – (H6'). Then for all $x \in \mathfrak{X}$, $T \in \mathfrak{T}$, $z \in \mathcal{Z}$ and $\theta_1 \in \Theta$, for $k \geq 1$,

$$E_{\theta_0} [U_{x,T,z,\theta_j}]^k < \infty; j = 0, 1, \quad (6.3)$$

$$E_{\theta_0} [V_{x,T,z,\theta_1}]^k < \infty, \quad (6.4)$$

$$E_{\theta_0} [V_{x,T,z,\theta_0,\theta_j}]^k < \infty; j = 0, 1. \quad (6.5)$$

Moreover, the above expectations are continuous in (x, T, z, θ_1) .

In particular, the Kullback-Leibler distance is continuous in x, T, z and θ_1 . The following lemma asserts that the average of the Kullback-Leibler distance is also a Kullback-Leibler distance in the limit.

Lemma 11 The limiting average $\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \mathcal{K}(f_{\theta_0, x^k, T_k, z_k}, f_{\theta_1, x^k, T_k, z_k})$ is also a Kullback-Leibler distance.

Even in this non-*iid* context, the Bayes factor is of the same form as (2.6); however, for $j = 0, 1$, $U_{x^i, T_i, z_i, \beta_j, \xi_j}$ and $V_{x^i, T_i, z_i, \beta_j, \xi_j}$ are not identically distributed for $i = 1, \dots, n$. Next, we establish strong consistency of Bayes factor in the non-*iid* SDE set-up by verifying the sufficient conditions of Theorem 7.

6.1 Verification of (5.2) and the Kullback-Leibler property in the non-*iid* set-up

Firstly, note that in our case,

$$\mathcal{K}^\infty(f_0, f) = \mathcal{K}^\infty(f_{\theta_0}, f_{\theta_1}), \quad (6.6)$$

where the rightmost side, as asserted by Lemma 11, clearly exists almost surely with respect to θ_1 and is also continuous in θ_1 .

Now note that compactness of $\mathfrak{X}$, $\mathfrak{T}$ and $\mathcal{Z}$ along with continuity of the function ϕ_{ξ_j} and $\mathcal{K}(f_{\theta_0, x, T, z}, f_{\theta_1, x, T, z})$ with respect to x, T and z implies

$$\begin{aligned} \psi(\theta_1) &= \inf_{x \in \mathfrak{X}, T \in \mathfrak{T}, z \in \mathcal{Z}} \mathcal{K}(f_{\theta_0, x, T, z}, f_{\theta_1, x, T, z}) \\ &= \inf_{x \in \mathfrak{X}, T \in \mathfrak{T}, z \in \mathcal{Z}} \int_0^T \left\{ \frac{\phi_{\xi_0}^2(z(s))}{2} E_{\theta_0}(\check{V}_{x, \beta_0}(s)) \right. \\ &\quad \left. - \phi_{\xi_0}(z(s)) \phi_{\xi_1}(z(s)) E_{\theta_0}(\check{V}_{x, \beta_0, \beta_1}(s)) + \frac{\phi_{\xi_1}^2(z(s))}{2} E_{\theta_0}(\check{V}_{x, \beta_1}(s)) \right\} ds \\ &= \inf_{x \in \mathfrak{X}, T \in \mathfrak{T}, z(\tilde{s}(T)) \in \mathcal{Z}} T \left\{ \frac{\phi_{\xi_0}^2(z(\tilde{s}(T)))}{2} E_{\theta_0}(\check{V}_{x, \beta_0}(\tilde{s}(T))) \right. \\ &\quad \left. - \phi_{\xi_0}(z(\tilde{s}(T))) \phi_{\xi_1}(z(\tilde{s}(T))) E_{\theta_0}(\check{V}_{x, \beta_0, \beta_1}(\tilde{s}(T))) + \frac{\phi_{\xi_1}^2(z(\tilde{s}(T)))}{2} E_{\theta_0}(\check{V}_{x, \beta_1}(\tilde{s}(T))) \right\}, \end{aligned} \quad (6.7)$$

by the mean value theorem for integrals, where $\tilde{s}(T) \in [0, T]$ such that the above equality holds. Also note that the expression in (6.7) is continuous in T since originally the integral on $[0, T]$ is continuous in T . Now note that if $|T - \tilde{T}| < \delta_1(\epsilon)$ such that $|\phi_{\xi_j}(z(\tilde{s}(T))) - \phi_{\xi_j}(z(\tilde{s}(\tilde{T})))| < \frac{\epsilon}{2}$ due to continuity in T and if $|z(\tilde{s}(\tilde{T})) - \tilde{z}(\tilde{s}(\tilde{T}))| < \delta_2(\epsilon)$ such that $|\phi_{\xi_j}(z(\tilde{s}(\tilde{T}))) - \phi_{\xi_j}(\tilde{z}(\tilde{s}(\tilde{T})))| < \frac{\epsilon}{2}$ due to continuity of ϕ_{ξ_j} in z , then $|\phi_{\xi_j}(z(\tilde{s}(T))) - \phi_{\xi_j}(\tilde{z}(\tilde{s}(\tilde{T})))| \leq |\phi_{\xi_j}(z(\tilde{s}(T))) - \phi_{\xi_j}(z(\tilde{s}(\tilde{T})))| + |\phi_{\xi_j}(z(\tilde{s}(\tilde{T}))) - \phi_{\xi_j}(\tilde{z}(\tilde{s}(\tilde{T})))| < \epsilon$, showing that $\phi_{\xi_j}(z(\tilde{s}(T)))$ is continuous in T and $z(\tilde{s}(T))$, which also belong to

compact spaces. Hence, from (6.7) it follows that

$$\begin{aligned} \psi(\theta_1) = T^*(\theta_1) & \left[\frac{\phi_{\xi_0}^2(z^*(\theta_1)(\tilde{s}(T^*(\theta_1))))}{2} E_{\theta_0}(\check{V}_{x^*(\theta_1), \beta_0}(\tilde{s}(T^*(\theta_1)))) \right. \\ & - \phi_{\xi_0}(z^*(\theta_1)(\tilde{s}(T^*(\theta_1)))) \phi_{\xi_1}(z^*(\theta_1)(\tilde{s}(T^*(\theta_1)))) E_{\theta_0}(\check{V}_{x^*(\theta_1), \beta_0, \beta_1}(\tilde{s}(T^*(\theta_1)))) \\ & \left. + \frac{\phi_{\xi_1}^2(z^*(\theta_1)(\tilde{s}(T^*(\theta_1))))}{2} E_{\theta_0}(\check{V}_{x^*(\theta_1), \beta_1}(\tilde{s}(T^*(\theta_1)))) \right], \end{aligned} \quad (6.8)$$

where $x^*(\theta_1) \in \mathfrak{X}$, $T^*(\theta_1) \in \mathfrak{T}$, $z^*(\theta_1)(\tilde{s}(T^*(\theta_1))) \in \mathfrak{Z}$ depend upon θ_1 . Then, considering the constant correspondence function $\gamma(\theta_1) = \mathfrak{X} \times \mathfrak{T} \times \mathfrak{Z}$, for all $\theta_1 \in \Theta$, we note that γ is both upper and lower hemicontinuous (hence continuous), and also compact-valued. Hence, Berge's maximum theorem (Berge (1963)) guarantees that (6.8) is a continuous function of θ_1 .

Because of continuity of $\psi(\theta_1)$ in θ_1 , the set $\{\theta_1 : \psi(\theta_1) \geq \delta\}$ is open and can be assigned any desired probability by choosing appropriate priors dominated by the Lebesgue measure. That is, we can assign prior probability one to this set by choosing appropriate priors dominated by the Lebesgue measure. Now, because of the inequality

$$\pi \left(\theta_1 : \inf_i \mathcal{K}(f_{\theta_0, x^i, T_i, z_i}, f_{\theta_1, x^i, T_i, z_i}) \geq \delta \right) \geq \pi(\theta_1 : \psi(\theta_1) \geq \delta),$$

and since we choose π such that $\pi(\theta_1 : \psi(\theta_1) \geq \delta) = 1$, it follows that

$$\pi \left(\theta_1 : \inf_i \mathcal{K}(f_{\theta_0, x^i, T_i, z_i}, f_{\theta_1, x^i, T_i, z_i}) \geq \delta \right) = 1,$$

satisfying (5.2).

The Kullback-Leibler property of the Lebesgue measure dominated π easily follows from continuity of $\mathcal{K}^\infty(f_{\theta_0}, f_{\theta_1})$ in θ_1 .

6.2 Verification of the Q^* property in the non-*iid* set-up

Observe that in this situation, for any $\epsilon > 0$,

$$\begin{aligned} A_n(\epsilon) &= \{f_n : \mathcal{K}(f_{0n}, f_n) \geq \epsilon\} \\ &= \{\theta_1 : \mathcal{K}(f_{\theta_0, x^n, T_n, z_n}, f_{\theta_1, x^n, T_n, z_n}) \geq \epsilon\} \end{aligned}$$

Then note that

$$\hat{f}_{nA_n(\epsilon)}(X) \leq \sup_{\theta_1 \in A_n(\epsilon)} f_{\theta_1, x^n, T_n, z_n}(X) = f_{\hat{\theta}_1(X, x^n, T_n, z_n)}(X), \quad (6.9)$$

where $\hat{\theta}_1(X, x^n, T_n, \mathbf{z}_n)$, which depends upon $X, x^n, T_n, \mathbf{z}_n$, is the maximizer lying in the compact set $A_n(\epsilon)$. Now,

$$\begin{aligned}
\mathcal{K}(f_{\theta_0, x^n, T_n, \mathbf{z}_n}, \hat{f}_{nA_n(\epsilon)}) &= E_{\theta_0} [\log f_{\theta_0, x^n, T_n, \mathbf{z}_n}(X)] - E_{\theta_0} [\log \hat{f}_{nA_n(\epsilon)}(X)] \\
&\geq E_{\theta_0} [\log f_{\theta_0, x^n, T_n, \mathbf{z}_n}(X)] - E_{\theta_0} [\log f_{\hat{\theta}_1(X, x^n, T_n, \mathbf{z}_n)}(X)] \\
&= E_{\theta_0} \left(\log \frac{f_{\theta_0, x^n, T_n, \mathbf{z}_n}(X)}{f_{\hat{\theta}_1(X, x^n, T_n, \mathbf{z}_n)}(X)} \right) \\
&= E_{\hat{\theta}_1(X, x^n, T_n, \mathbf{z}_n) | \theta_0} E_{X | \hat{\theta}_1(X, x^n, T_n, \mathbf{z}_n) = \varphi, \theta_0} \left(\log \frac{f_{\theta_0, x^n, T_n, \mathbf{z}_n}(X)}{f_{\{\hat{\theta}_1(X, x^n, T_n, \mathbf{z}_n) = \varphi\}}(X)} \right) \\
&= E_{\hat{\theta}_1(X, x^n, T_n, \mathbf{z}_n) | \theta_0} \mathcal{K}(f_{\theta_0, x^n, T_n, \mathbf{z}_n}, f_{\varphi, x^n, T_n, \mathbf{z}_n}) \\
&\geq E_{\hat{\theta}_1(X, x^n, T_n, \mathbf{z}_n) | \theta_0} \inf_{\boldsymbol{\eta} \in A_n(\epsilon)} \mathcal{K}(f_{\theta_0, x^n, T_n, \mathbf{z}_n}, f_{\boldsymbol{\eta}, x^n, T_n, \mathbf{z}_n}) \\
&= E_{\hat{\theta}_1(X, x^n, T_n, \mathbf{z}_n) | \theta_0} \mathcal{K}(f_{\theta_0, x^n, T_n, \mathbf{z}_n}, f_{\boldsymbol{\eta}_n^*, x^n, T_n, \mathbf{z}_n}) \\
&= \mathcal{K}(f_{\theta_0, x^n, T_n, \mathbf{z}_n}, f_{\boldsymbol{\eta}_n^*, x^n, T_n, \mathbf{z}_n}) \\
&\geq \epsilon,
\end{aligned} \tag{6.10}$$

where $\boldsymbol{\eta}_n^* = \arg \min_{\boldsymbol{\eta} \in A_n(\epsilon)} \mathcal{K}(f_{\theta_0, x^n, T_n, \mathbf{z}_n}, f_{\boldsymbol{\eta}, x^n, T_n, \mathbf{z}_n}) \in A_n(\epsilon)$, due to compactness of $A_n(\epsilon)$. Hence, (5.4) is satisfied in our non-*iid* SDE set-up.

6.3 Verification of (5.7)

From Lemma 10 it follows that $E \left\{ \log \frac{f_{\theta_0, x, T, \mathbf{z}}(X)}{f_{\theta_1, x, T, \mathbf{z}}(X)} \right\}^2$ exists and is continuous in θ_1, x, T and $\mathbf{z}$. Then compactness of $\Theta, \mathfrak{X}, \mathfrak{T}$ and $\mathcal{Z}$ ensures (5.7).

6.4 Verification of (5.8)

For the non-*iid* case, the following identity holds:

$$\begin{aligned}
\frac{I_{n+1}}{I_n} &= \frac{\hat{f}_{n+1}(X_{n+1})}{f_{0, n+1}(X_{n+1})} \\
&= \frac{\hat{f}_{x^{n+1}, T_{n+1}, \mathbf{z}_{n+1}}(X_{n+1})}{f_{\theta_0, x^{n+1}, T_{n+1}, \mathbf{z}_{n+1}}(X_{n+1})},
\end{aligned} \tag{6.11}$$

where

$$\hat{f}_{x^{n+1}, T_{n+1}, \mathbf{z}_{n+1}}(\cdot) = E_{\theta_1 | X_1, \dots, X_n} [f_{\theta_1, x^{n+1}, T_{n+1}, \mathbf{z}_{n+1}}(\cdot)] \tag{6.12}$$

is the posterior predictive distribution of $f_{\theta_1, x^{n+1}, T_{n+1}, \mathbf{z}_{n+1}}(\cdot)$, with respect to the posterior of θ_1 , given $X_1, \dots, X_n$.

Now since $\log f_{\theta_0, x^{n+1}, T_{n+1}, \mathbf{z}_{n+1}}(X_{n+1}) = U_{x^{n+1}, T_{n+1}, \mathbf{z}_{n+1}, \theta_0} - \frac{V_{x^{n+1}, T_{n+1}, \mathbf{z}_{n+1}, \theta_0}}{2}$, using Lemma 10 and compactness of $\Theta, \mathfrak{X}, \mathfrak{T}$ and $\mathcal{Z}$ it is easy to see that the moments of $\log f_{\theta_0, x^{n+1}, T_{n+1}, \mathbf{z}_{n+1}}(X_{n+1})$ are uniformly bounded above. So, we have

$$\sup_n E \left(\log f_{\theta_0, x^{n+1}, T_{n+1}, \mathbf{z}_{n+1}}(X_{n+1}) \right)^2 < \infty. \tag{6.13}$$

As in the *iid* case, here also we have

$$\begin{aligned} f_{\theta_1^*}(X_{n+1}, x^{n+1}, T_{n+1}, \mathbf{z}_{n+1})(X_{n+1}) &= \inf_{\theta_1 \in \Theta} f_{\theta_1, x^{n+1}, T_{n+1}, \mathbf{z}_{n+1}}(X_{n+1}) \\ &\leq \hat{f}_{x^{n+1}, T_{n+1}, \mathbf{z}_{n+1}}(X_{n+1}) \\ &\leq \sup_{\theta_1 \in \Theta} f_{\theta_1, x^{n+1}, T_{n+1}, \mathbf{z}_{n+1}}(X_{n+1}) = f_{\theta_1^{**}}(X_{n+1}, x^{n+1}, T_{n+1}, \mathbf{z}_{n+1})(X_{n+1}), \end{aligned}$$

where $\theta_1^*(X_{n+1}, x^{n+1}, T_{n+1}, \mathbf{z}_{n+1}) = \arg \min_{\theta_1 \in \Theta} f_{\theta_1, x^{n+1}, T_{n+1}, \mathbf{z}_{n+1}}(X_{n+1}) \in \Theta$ and $\theta_1^{**}(X_{n+1}, x^{n+1}, T_{n+1}, \mathbf{z}_{n+1}) = \arg \max_{\theta_1 \in \Theta} f_{\theta_1, x^{n+1}, T_{n+1}, \mathbf{z}_{n+1}}(X_{n+1}) \in \Theta$. Note that each component of $\theta_1^*(X_{n+1}, x^{n+1}, T_{n+1}, \mathbf{z}_{n+1})$ and $\theta_1^{**}(X_{n+1}, x^{n+1}, T_{n+1}, \mathbf{z}_{n+1})$ depends on $X_{n+1}, x^{n+1}, T_{n+1}, \mathbf{z}_{n+1}$.

It follows, as in the *iid* case, that

$$\begin{aligned} & - \left| U_{\theta_1^*, x^{n+1}, T_{n+1}, \mathbf{z}_{n+1}} \right| - \frac{V_{\theta_1^*, x^{n+1}, T_{n+1}, \mathbf{z}_{n+1}}}{2} \\ & \leq U_{\theta_1^*, x^{n+1}, T_{n+1}, \mathbf{z}_{n+1}} - \frac{V_{\theta_1^*, x^{n+1}, T_{n+1}, \mathbf{z}_{n+1}}}{2} \\ & \leq \log \hat{f}_{x^{n+1}, T_{n+1}, \mathbf{z}_{n+1}}(X_{n+1}) \\ & \leq U_{\theta_1^{**}, x^{n+1}, T_{n+1}, \mathbf{z}_{n+1}} - \frac{V_{\theta_1^{**}, x^{n+1}, T_{n+1}, \mathbf{z}_{n+1}}}{2} \\ & \leq \left| U_{\theta_1^{**}, x^{n+1}, T_{n+1}, \mathbf{z}_{n+1}} \right| + \frac{V_{\theta_1^{**}, x^{n+1}, T_{n+1}, \mathbf{z}_{n+1}}}{2}. \end{aligned}$$

Proceeding in the same way as in the *iid* case, and exploiting Lemma 10, we obtain

$$\sup_n E \left(\log \hat{f}_{x^{n+1}, T_{n+1}, \mathbf{z}_{n+1}}(X_{n+1}) \right)^2 < \infty. \quad (6.14)$$

Thus, as in the *iid* set-up, (5.8) follows from (6.13) and (6.14).

We formalize the above arguments in the form of a theorem in our non-*iid* SDE set-up.

Theorem 12 Assume the non-*iid* SDE set-up and conditions (H1') – (H6'). Then (5.9) holds.

As in the previous cases, the following corollary provides asymptotic comparison between two models using Bayes factor in the non-*iid* SDE set-up.

Corollary 13 For $j = 1, 2$, let $R_{jn}(\theta_j) = \prod_{i=1}^n \frac{f_{\theta_j, x^i, T_i, \mathbf{z}_i}(X_i)}{f_{\theta_0, x^i, T_i, \mathbf{z}_i}(X_i)}$, where θ_1 and θ_2 are two different finite sets of parameters, perhaps with different dimensionalities, associated with the two models to be compared. For $j = 1, 2$, let

$$I_{jn} = \int R_{jn}(\theta_j) \pi_j(d\theta_j),$$

where π_j is the prior on θ_j . Let $B_n = I_{1n}/I_{2n}$ as before. Assume the non-*iid* SDE set-up and suppose that both the models satisfy (H1') – (H6'), and have the Kullback-Leibler property with $\delta = \delta_1$ and $\delta = \delta_2$ respectively. Then

$$n^{-1} \log B_n \rightarrow \delta_2 - \delta_1,$$

almost surely.

7 Simulation studies

7.1 Covariate selection when $n = 15, T = 1$

We demonstrate with simulation study the finite sample analogue of Bayes factor analysis as $n \rightarrow \infty$ and T is fixed. In this regard, we consider $n = 15$ individuals, where the i -th one is modeled by

$$dX_i(t) = (\xi_1 + \xi_2 z_1(t) + \xi_3 z_2(t) + \xi_4 z_3(t))(\xi_5 + \xi_6 X_i(t))dt + \sigma_i dW_i(t), \quad (7.1)$$

for $i = 1, \dots, 15$. We fix our diffusion coefficients as $\sigma_{i+1} = \sigma_i + 5$ for $i = 1, \dots, 14$ where $\sigma_1 = 10$. We consider the initial value $X(0) = 0$ and the time interval $[0, T]$ with $T = 1$.

To achieve numerical stability of the marginal likelihood corresponding to each data we choose the true values of $\xi_i; i = 1, \dots, 6$ as follows: $\xi_i \stackrel{iid}{\sim} N(\mu_i, 0.001^2)$, where $\mu_i \stackrel{iid}{\sim} N(0, 1)$. This is not to be interpreted as the prior; this is just a means to set the true values of the parameters of the data-generating model.

We assume that the time dependent covariates $z_i(t)$ satisfy the following *SDEs*

$$\begin{aligned} dz_1(t) &= (\tilde{\theta}_1 + \tilde{\theta}_2 z_1(t))dt + dW_1(t) \\ dz_2(t) &= \tilde{\theta}_3 dt + dW_2(t) \\ dz_3(t) &= \tilde{\theta}_4 z_3(t)dt + dW_3(t), \end{aligned} \quad (7.2)$$

where $W_i(\cdot); i = 1, 2, 3$, are independent Wiener processes, and $\tilde{\theta}_i \stackrel{iid}{\sim} N(0, 0.01^2)$ for $i = 1, \dots, 4$.

We obtain the covariates by first simulating $\tilde{\theta}_i \stackrel{iid}{\sim} N(0, 0.01^2)$ for $i = 1, \dots, 4$, fixing the values, and then by simulating the covariates using the *SDEs* (7.2) by discretizing the time interval $[0, 1]$ into 500 equispaced time points. In all our applications we have standardized the covariates over time so that they have zero means and unit variances.

Once the covariates are thus obtained, we assume that the data are generated from the (true) model where all the covariates are present. For the true values of the parameters, we simulated $(\xi_1, \dots, \xi_6)$ from the prior and treated the obtained values as the true set of parameters θ_0 . We then generated the data using (7.1) by discretizing the time interval $[0, 1]$ into 500 equispaced time points.

As we have three covariates so we will have $2^3 = 8$ different models. Denoting a model by the presence and absence of the respective covariates, it then is the case that $(1, 1, 1)$ is the true, data-generating model, while $(0, 0, 0)$, $(0, 0, 1)$, $(0, 1, 0)$, $(0, 1, 1)$, $(1, 0, 0)$, $(1, 0, 1)$, and $(1, 1, 0)$ are the other 7 possible models.

7.1.1 Case 1: the true parameter set θ_0 is fixed

Prior on θ

For the prior π on θ , we first obtain the maximum likelihood estimator (*MLE*) of θ using simulated annealing (see, for example, Liu (2001), Robert and Casella (2004)), and consider a normal prior where the mean is the *MLE* of ξ_i for $i = 1, \dots, 6$ and the variance is $0.8^2 \mathbb{I}_6$, $\mathbb{I}_6$ being the 6-dimensional identity matrix. As will be seen, this results in consistent model selection using Bayes factor.

Form of the Bayes factor

In this case the related Bayes factor has the form

$$I_n = \int \prod_{i=1}^n \frac{f_{i, \theta_1}(X_i)}{f_{i, \theta_0}(X_i)} \pi(d\theta_1),$$

where $\theta_0 = (\xi_{0,1}, \xi_{0,2}, \xi_{0,3}, \xi_{0,4}, \xi_{0,5}, \xi_{0,6})$ is the true parameter set and $\theta_1 = (\xi_1, \xi_2, \xi_3, \xi_4, \xi_5, \xi_6)$ is the unknown set of parameters corresponding to any other model. Table 7.1 describes the results of our

Bayes factor analyses. It is clear from the 7 values of the table that the correct model $(1, 1, 1)$ is always

Table 7.1: Bayes factor results

Model	$\frac{1}{15} \log I_{15}$
(0, 0, 0)	-3.25214
(0, 0, 1)	-1.39209
(0, 1, 0)	-3.31954
(0, 1, 1)	-1.11729
(1, 0, 0)	-3.40378
(1, 0, 1)	-1.22529
(1, 1, 0)	-3.46790

preferred.

7.1.2 Case 2: the parameter set θ_0 is random and has the prior distribution π

We consider the same form of the prior π as in Section 7.1.1, but with variance $0.1^2 \mathbb{I}_6$. The smaller variance compared to that in Case 1 attempts to somewhat compensate, in essence, for the lack of precise information about the true parameter values.

In this case we calculate the marginal log-likelihood of the 8 possible models as

$$\ell_i = \frac{1}{15} \log \int \prod_{i=1}^n f_{i,\theta_1}(X_i) \pi(d\theta_1); \quad i = 1, \dots, 8,$$

with ℓ_8 corresponding to the true model. Table 7.2 shows that ℓ_8 is the highest. This clearly implies that the Bayes factor consistently selects the correct set of covariates even though the parameters of the true model are not fixed.

Table 7.2: Values of $\frac{1}{15} \times$ marginal log-likelihoods

Model	ℓ_i
(0, 0, 0)	2.42430
(0, 0, 1)	4.29608
(0, 1, 0)	1.75213
(0, 1, 1)	4.84717
(1, 0, 0)	1.56242
(1, 0, 1)	4.92628
(1, 1, 0)	0.47111
(1, 1, 1)	5.84665 (true model)

8 Summary and conclusion

In this article we have investigated the asymptotic theory of Bayes factors when the models are associated with systems of *SDE*'s consisting of sets of time-dependent covariates. The model selection problem we consider encompasses appropriate selection of a subset of covariates, as well as appropriate selection of the part of the drift function that does not involve covariates. Such an undertaking, according to our knowledge, is a first-time effort which did not hitherto take place in the literature.

We have established almost sure exponential convergence of the Bayes factor when the time domains remain bounded but the number of individuals tend to infinity, in both *iid* and non-*iid* cases. In the non-*iid* context, we proposed and proved general results on Bayes factor asymptotics, which should be of independent interest.

Our simulation studies demonstrate that Bayes factor is a reliable criterion even in non-asymptotic situations for capturing the correct set of covariates in our *SDE* set-ups.

Note that our theory for non-*iid* situations readily extends to model comparison problems when one of the models is associated with an *iid* system of *SDE*'s and another with a non-*iid* system of *SDE*'s. For instance, if the true model is associated with an *iid* system, then $f_{0i} \equiv f_0 \equiv f_{\theta_0}$, and the rest of the theory remains the same as our non-*iid* theory of Bayes factors. The case when the other model is associated with an *iid* system is analogous.

Acknowledgments

The first author gratefully acknowledges her CSIR Fellowship, Govt. of India.

Supplementary Material

Throughout, we refer to our main manuscript Maitra and Bhattacharya (2016a) as MB.

S-1 Proof of Lemma 1 of MB

We first consider $k = 1$. Note that due to assumption (H4'),

$$E_{\theta_0}(V_{i,\theta_1}) \leq T_i K_{\beta_1} \left(\sup_{s \in [0, T_i]} \phi_{i,\xi_j}^2(s) \right) \left(1 + \sup_{s \in [0, T_i]} E_{\theta_0}(X_i^2(s)) + \|\beta_1\|^2 \right) < \infty,$$

since, by Proposition 1 of Delattre *et al.* (2013), $\sup_{s \in [0, T_i]} E_{\theta_0}(X_i^{2\ell}(s)) < \infty$, for $\ell \geq 1$, and since

$\left(\sup_{s \in [0, T_i]} \phi_{i,\xi_j}^2(s) \right)$ is bounded above due to continuity of g_l ; $l = 1, \dots, p$. Hence, (3.6) of MB holds.

Now observe that due to Cauchy-Schwartz and (H4')

$$\begin{aligned} E_{\theta_0}(V_{i,\theta_0,\theta_j}) &= \int_0^{T_i} E_{\theta_0} \left(\frac{\phi_{i,\xi_0}(s) \phi_{i,\xi_j}(s) b_{\beta_j}(X_i(s)) b_{\beta_0}(X_i(s))}{\sigma^2(X_i(s))} \right) ds \\ &\leq \int_0^{T_i} \left[E_{\theta_0} \left(\frac{\phi_{i,\xi_j}^2(s) b_{\beta_j}^2(X_i(s))}{\sigma^2(X_i(s))} \right) \right]^{\frac{1}{2}} \times \left[E_{\theta_0} \left(\frac{\phi_{i,\xi_0}^2(s) b_{\beta_0}^2(X_i(s))}{\sigma^2(X_i(s))} \right) \right]^{\frac{1}{2}} ds \\ &\leq T_i \left(\sup_{s \in [0, T_i]} \phi_{i,\xi_j}^2(s) \right)^{\frac{1}{2}} K_{\beta_j}^{\frac{1}{2}} \left(1 + \sup_{s \in [0, T_i]} E_{\theta_0}(X_i^2(s)) + \|\beta_j\|^2 \right)^{\frac{1}{2}} \\ &\quad \times K_{\beta_0}^{\frac{1}{2}} \left(\sup_{s \in [0, T_i]} \phi_{i,\xi_0}^2(s) \right)^{\frac{1}{2}} \left(1 + \sup_{s \in [0, T_i]} E_{\theta_0}(X_i^2(s)) + \|\beta_0\|^2 \right)^{\frac{1}{2}} \\ &< \infty, \end{aligned}$$

by Proposition 1 of Delattre *et al.* (2013) and boundedness of $\left(\sup_{s \in [0, T_i]} \phi_{i,\xi_j}^2(s) \right)$. Hence, (3.7) of MB holds. Also note that since $E_{\theta_0}(U_{i,\theta_j}) = E_{\theta_0}(V_{i,\theta_0,\theta_j})$ by (2.11) and (2.13), (3.5) is implied by (3.7) of MB. To see that the moments are continuous in θ_1 , let $\{\theta_1^{(m)}\}_{m=1}^{\infty}$ be a sequence converging to $\tilde{\theta}_1$ as $m \rightarrow \infty$. Due to (H3'),

$$\frac{\phi_{i,\xi_1}^2(s) b_{\beta_1}^2(X_i(s))}{\sigma^2(X_i(s))} \rightarrow \frac{\phi_{i,\tilde{\xi}_1}^2(s) b_{\tilde{\beta}_1}^2(X_i(s))}{\sigma^2(X_i(s))}$$

and

$$\frac{\phi_{i,\xi_1}^{(m)}(s) \phi_{i,\xi_0}(s) b_{\beta_1}^{(m)}(X_i(s)) b_{\beta_0}(X_i(s))}{\sigma^2(X_i(s))} \rightarrow \frac{\phi_{i,\tilde{\xi}_1}(s) \phi_{i,\xi_0}(s) b_{\tilde{\beta}_1}(X_i(s)) b_{\beta_0}(X_i(s))}{\sigma^2(X_i(s))},$$

as $m \rightarrow \infty$, for any given sample path $\{X_i(s) : s \in [0, T_i]\}$. Assumption (H4') implies that $\frac{\phi_{i,\xi_1}^2(s) b_{\beta_1}^2(X_i(s))}{\sigma^2(X_i(s))}$

is dominated by $\sup_{\xi_1 \in \Gamma, s \in [0, T_i]} \phi_{i,\xi_1}^2(s) \times \sup_{\beta_1 \in \mathfrak{B}} K_{\beta_1} \left(1 + \sup_{s \in [0, T_i]} [X_i(s)]^2 + \sup_{\beta_1 \in \mathfrak{B}} \|\beta_1\|^2 \right)$. Since $X_i(s)$ is continuous on $[0, T_i]$, (guaranteed by (H2')); see Delattre *et al.* (2013)), it follows that $\int_0^{T_i} [X_i(s)]^2 ds <$

∞ , which, in turn guarantees, in conjunction with compactness of $\mathfrak{B}$ and Γ , that the upper bound is integrable. Hence, $V_{i,\theta_1^{(m)}} \rightarrow V_{i,\tilde{\theta}_1}$, almost surely. Now, for all $m \geq 1$,

$$V_{i,\theta_1^{(m)}} < T_i \left(\sup_{\xi_1 \in \Gamma, s \in [0, T_i]} \phi_{i,\xi_1}^2(s) \right) \times \left(\sup_{\beta_1 \in \mathfrak{B}} K_{\beta_1} \right) \times \left(1 + \sup_{s \in [0, T_i]} [X_i(s)]^2 + \sup_{\beta_1 \in \mathfrak{B}} \|\beta_1\|^2 \right).$$

Since $E_{\theta_0} \left(\sup_{s \in [0, T_i]} [X_i(s)]^2 \right) < \infty$ by (3.1) of MB, it follows that $E_{\theta_0} (V_{i,\theta_1^{(m)}}) \rightarrow E_{\theta_0} (V_{i,\tilde{\theta}_1})$, as $(\theta_1^{(m)}) \rightarrow (\tilde{\theta}_1)$. Hence, $E_{\theta_0} (V_{i,\theta_1})$ is continuous in θ_1 .

In the case of V_{i,θ_0,θ_1} , the relevant quantity $\frac{\phi_{i,\xi_1^{(m)}} \phi_{i,\xi_0} b_{\beta_1^{(m)}}(X_i(s)) b_{\beta_0}(X_i(s))}{\sigma^2(X_i(s))}$ is dominated by the continuous function (hence integrable on $[0, T_i]$)

$$\begin{aligned} & \left(\sup_{\xi_1 \in \Gamma, s \in [0, T_i]} |\phi_{i,\xi_1}(s)| \right) \times \left(\sup_{\beta_1 \in \mathfrak{B}} K_{\beta_1}^{\frac{1}{2}} \right) \times \left(1 + \sup_{s \in [0, T_i]} [X_i(s)]^2 + \sup_{\beta_1 \in \mathfrak{B}} \|\beta_1\|^2 \right) \\ & \times \left(\sup_{s \in [0, T_i]} |\phi_{i,\xi_0}(s)| \right) \times K_{\beta_0}^{\frac{1}{2}} \left(1 + \sup_{s \in [0, T_i]} [X_i(s)]^2 + \|\beta_0\|^2 \right), \end{aligned}$$

which ensures $V_{i,\theta_0,\theta_1^{(m)}} \rightarrow V_{i,\theta_0,\tilde{\theta}_1}$, almost surely. Using the above bound for

$\frac{\phi_{i,\xi_1^{(m)}} \phi_{i,\xi_0} b_{\beta_1^{(m)}}(X_i(s)) b_{\beta_0}(X_i(s))}{\sigma^2(X_i(s))}$, it is seen that

$$V_{i,\theta_0,\theta_1^{(m)}} < T_i K_1 \left(\sup_{s \in [0, T_i]} [X_i(s)]^4 + K_2 \sup_{s \in [0, T_i]} [X_i(s)]^2 + K_3 \right),$$

for appropriate positive constants K_1, K_2, K_3 , so that (3.1) of MB for $k = 4$, guarantees that $E_{\theta_0} (V_{i,\theta_0,\theta_1^{(m)}}) \rightarrow E_{\theta_0} (V_{i,\theta_0,\tilde{\theta}_1})$, as $\theta_1^{(m)} \rightarrow \tilde{\theta}_1$. This shows that $E_{\theta_0} (V_{i,\theta_0,\theta_1})$ is continuous as well. Since $E_{\theta_0} (U_{i,\theta_1}) = E_{\theta_0} (V_{i,\theta_0,\theta_1})$, it follows that $E_{\theta_0} (U_{i,\theta_1})$ is continuous in θ_1 .

We now consider $k \geq 2$. Note that, due to (H4'), and the inequality $(a+b)^k \leq 2^{k-1}(|a|^k + |b|^k)$ for $k \geq 2$ and any a, b ,

$$\begin{aligned} E_{\theta_0} (V_{i,\theta_j})^k & \leq \left(\sup_{s \in [0, T_i]} |\phi_{i,\xi_j}(s)|^k \right) 2^{k-1} T_i^k K_{\beta_j}^k (1 + \|\beta_j\|^2)^k \\ & \quad + \left(\sup_{s \in [0, T_i]} |\phi_{i,\xi_j}(s)|^k \right) 2^{k-1} T_i^k K_{\beta_j}^k E \left(\sup_{s \in [0, T_i]} [X_i(s)]^{2k} \right). \end{aligned}$$

Since $E \left(\sup_{s \in [0, T_i]} [X_i(s)]^{2k} \right) < \infty$ due to (3.1) of MB, and because $K_{\beta_j}, \|\beta_j\|$ are continuous in compact $\mathfrak{B}$, and $\left(\sup_{s \in [0, T_i]} |\phi_{i,\xi_j}(s)|^k \right)$ is continuous in compact Γ , it holds that $E_{\theta_0} (V_{i,\theta_j})^k < \infty$. In a similar manner it can be shown that $E_{\theta_0} (V_{i,\theta_0,\theta_1})^k < \infty$. Thus, (3.7) of MB follows.

To see that (3.5) of MB holds, note that, due to (2.11) of MB and $(a+b)^k \leq 2^{k-1}(|a|^k + |b|^k)$,

$$E_{\theta_0} (U_{i,\theta_j})^k \leq 2^{k-1} E_{\theta_0} (V_{i,\theta_0,\theta_j})^k + 2^{k-1} E_{\theta_0} \left(\int_0^{T_i} \frac{\phi_{i,\xi_j}(s) b_{\beta_j}(X_i(s))}{\sigma(X_i(s))} dW_i(s) \right)^k. \quad (\text{S-1.1})$$

Since, due to (H4'), (3.1) of MB and continuity of ϕ_{i,ξ_j} on compact spaces,

$$E_{\theta_0} \left(\int_0^T \left| \frac{\phi_{i,\xi_j}(s)b_{\beta_j}(X_i(s))}{\sigma(X_i(s))} \right|^k ds \right) < \infty,$$

Theorem 7.1 of Mao (2011) (page 39) shows that

$$E_{\theta_0} \left(\left| \int_0^{T_i} \frac{\phi_{i,\xi_j}(s)b_{\beta_j}(X_i(s))}{\sigma(X_i(s))} dW(s) \right|^k \right) \leq \left(\frac{k(k-1)}{2} \right)^{\frac{k}{2}} T_i^{\frac{k-2}{2}} E_{\theta_0} \left(\int_0^{T_i} \left| \frac{\phi_{i,\xi_j}(s)b_{\beta_j}(X_i(s))}{\sigma(X_i(s))} \right|^k ds \right). \quad (\text{S-1.2})$$

Combining (S-1.1) with (S-1.2) and the result $E_{\theta_0} (V_{i,\theta_0,\theta_1})^k < \infty$, it follows that $E_{\theta_0} (U_{i,\theta_j})^k < \infty$.

As regards continuity of the moments for $k \geq 2$, first note that in the context of $k = 1$, we have shown almost sure continuity of V_{i,θ_1} with respect to θ_1 . Hence, V_{i,θ_1}^k is almost surely continuous with respect to θ_1 . That is, $\theta_1^{(m)} \rightarrow \tilde{\theta}_1$ implies $V_{i,\theta_1^{(m)}}^k \rightarrow V_{i,\tilde{\theta}_1}^k$, almost surely. Once again, dominated convergence theorem allows us to conclude that $E_{\theta_0} (V_{i,\theta_1^{(m)}})^k \rightarrow E_{\theta_0} (V_{i,\tilde{\theta}_1})^k$, implying continuity of $E_{\theta_0} (V_{i,\theta_1})^k$ with respect to θ_1 . Similarly, it is easy to see that $E_{\theta_0} (V_{i,\theta_0,\theta_1})^k$ is continuous with respect to θ_1 . To see continuity of $E_{\theta_0} (U_{i,\theta_1})^k$, first note that

$$E_{\theta_0} \left[\int_0^{T_i} \left(\frac{\phi_{i,\xi_1^{(m)}}(s)b_{\beta_1^{(m)}}(X_i(s))}{\sigma(X_i(s))} - \frac{\phi_{i,\tilde{\xi}_1}(s)b_{\tilde{\beta}_1}(X_i(s))}{\sigma(X_i(s))} \right)^2 ds \right] \rightarrow 0,$$

as $m \rightarrow \infty$. The result follows as before by first noting pointwise convergence, and then using (H4') and then (3.1) of MB, along with (H1') and boundedness of $\phi_{i,\xi_1^{(m)}}$. By Itô isometry it holds that

$$E_{\theta_0} \left[\int_0^{T_i} \frac{\phi_{i,\xi_1^{(m)}}(s)b_{\beta_1^{(m)}}(X_i(s))}{\sigma(X_i(s))} dW_i(s) - \int_0^{T_i} \frac{\phi_{i,\tilde{\xi}_1}(s)b_{\tilde{\beta}_1}(X_i(s))}{\sigma(X_i(s))} dW_i(s) \right]^2 \rightarrow 0.$$

Hence,

$$\int_0^{T_i} \frac{\phi_{i,\xi_1^{(m)}}(s)b_{\beta_1^{(m)}}(X_i(s))}{\sigma(X_i(s))} dW_i(s) \rightarrow \int_0^{T_i} \frac{\phi_{i,\tilde{\xi}_1}(s)b_{\tilde{\beta}_1}(X_i(s))}{\sigma(X_i(s))} dW_i(s)$$

in probability, as $m \rightarrow \infty$. Since $V_{i,\theta_0,\theta_1^{(m)}} \rightarrow V_{i,\theta_0,\tilde{\theta}_1}$ almost surely as $m \rightarrow \infty$, it follows from (2.11) of MB that $U_{i,\theta_1^{(m)}} \rightarrow U_{i,\tilde{\theta}_1}$ in probability, so that $U_{i,\theta_1^{(m)}}^k \rightarrow U_{i,\tilde{\theta}_1}^k$ in probability. Using (H4'), (3.1) of MB and (H1'), it is easily seen, using the same methods associated with (S-1.1) and (S-1.2), that $\sup_m E_{\theta_0} (U_{i,\theta_1^{(m)}})^{2k} < \infty$, proving that $\left\{ U_{i,\theta_1^{(m)}}^k \right\}_{m=1}^{\infty}$ is uniformly integrable. Hence, $E_{\theta_0} (U_{i,\theta_1^{(m)}})^k \rightarrow E_{\theta_0} (U_{i,\tilde{\theta}_1})^k$. In other words, $E_{\theta_0} (U_{i,\theta_1})^k$ is continuous in θ_1 .

S-2 Proof of Theorem 7 of MB

Let us consider the martingale sequence

$$S_N = \sum_{n=1}^N [\log(I_n/I_{n-1}) + \mathcal{K}(f_{0n}, \hat{f}_n)],$$

which is a martingale because $E[\log(I_n/I_{n-1})|X_1, X_2, \dots, X_{n-1}] = -\mathcal{K}(f_{0n}, \hat{f}_n)$. Using the above it can be verified that if (5.8) of MB holds, implying

$$\sum_{n=1}^{\infty} n^{-2} \text{Var} \left[\log \frac{I_n}{I_{n-1}} \right] < \infty,$$

then $S_N/N \rightarrow 0$ almost surely. Therefore

$$N^{-1} \log I_N + N^{-1} \sum_{n=1}^N \mathcal{K}(f_{0n}, \hat{f}_n) \rightarrow 0, \quad (\text{S-2.1})$$

almost surely, as $N \rightarrow \infty$.

Now consider $N^{-1} \sum_{i=1}^N \log \frac{f_{0i}(X_i)}{f_i(X_i)}$. If (5.7) of MB holds, implying

$$\sum_{i=1}^{\infty} i^{-2} \text{Var} \left[\log \frac{f_{0i}(X_i)}{f_i(X_i)} \right] < \infty,$$

then by Kolmogorov's strong law of large numbers in the independent but non-identical case,

$$\frac{1}{N} \sum_{i=1}^N \log \frac{f_{0i}(X_i)}{f_i(X_i)} \rightarrow \mathcal{K}^{\infty}(f_0, f)$$

almost surely, as $N \rightarrow \infty$. Let $\mathcal{N}_0(c) = \{f : \delta \leq \mathcal{K}^{\infty}(f_0, f) \leq \delta + c\}$, where $c > 0$. Now, note that,

$$\begin{aligned} I_N &= \int \frac{\prod_{i=1}^N f_i(X_i)}{\prod_{i=1}^N f_{0i}(X_i)} \pi(\tilde{d}f) \\ &\geq \int_{\mathcal{N}_0(c)} \exp \left(\sum_{i=1}^N \log \frac{f_i(X_i)}{f_{0i}(X_i)} \right) \pi(\tilde{d}f) \\ &= \int_{\mathcal{N}_0(c)} \exp \left(- \sum_{i=1}^N \log \frac{f_{0i}(X_i)}{f_i(X_i)} \right) \pi(\tilde{d}f). \end{aligned}$$

By Jensen's inequality,

$$\frac{1}{N} \log I_N \geq - \int_{\mathcal{N}_0(c)} \frac{1}{N} \left(\sum_{i=1}^N \log \frac{f_{0i}(X_i)}{f_i(X_i)} \right) \pi(\tilde{d}f) \quad (\text{S-2.2})$$

The integrand on the right hand side converges to $\mathcal{K}^{\infty}(f_0, f)$, pointwise for every f , given any sequence $\{X_i\}_{i=1}^{\infty}$ associated with the complement of some null set. Since, for all such sequences, uniform integrability of the integrand is guaranteed by (5.7) of MB, it follows that the right hand side of (S-2.2) converges to $-\int_{\mathcal{N}_0(c)} \mathcal{K}^{\infty}(f_0, f) \pi(\tilde{d}f)$ almost surely. Hence, almost surely,

$$\begin{aligned} \liminf_N N^{-1} \log I_N &\geq - \int_{\mathcal{N}_0(c)} \mathcal{K}^{\infty}(f_0, f) \pi(\tilde{d}f) \\ &\geq -(\delta + c) \pi(\mathcal{N}_0(c)) \\ &\geq -(\delta + c). \end{aligned}$$

Since $c > 0$ is arbitrary, it follows that

$$\liminf_N N^{-1} \log I_N \geq -\delta, \quad (\text{S-2.3})$$

almost surely. Now, due to (5.2) of MB it follows that $\mathcal{K}(f_{0n}, f_n) \geq \delta$ for all n with probability 1, so that $\mathcal{K}(f_{0n}, \hat{f}_n) = \mathcal{K}(f_{0n}, \hat{f}_{n, A_n(\delta)})$, where $A_n(\delta)$ is given by (5.5) of MB. By the Q^* property it implies that

$$\liminf_N N^{-1} \sum_{n=1}^N \mathcal{K}(f_{0n}, \hat{f}_n) \geq \delta.$$

Hence, it follows from (S-2.1) that

$$\limsup_N N^{-1} \log I_N \leq -\delta. \quad (\text{S-2.4})$$

Combining (S-2.3) and (S-2.4) it follows that

$$\lim_{N \rightarrow \infty} N^{-1} \log I_N = -\delta,$$

almost surely.

S-3 Proof of Lemma 10 of MB

The proofs of (6.3) – (6.5) of MB follow in the same way as the proofs of (3.5) – (3.7) of MB, using compactness of $\mathfrak{X}$, $\mathfrak{T}$ and $\mathfrak{Z}$ in addition to that of $\mathfrak{B}$ and Γ .

For the proofs of continuity of the moments, note that as in the *iid* case, uniform integrability is ensured by (H4'), (3.1) of MB and compactness of the sets $\mathfrak{B}$, Γ , $\mathfrak{X}$, $\mathfrak{T}$ and $\mathfrak{Z}$. The rest of the proof is almost the same as the proof of Theorem 5 of Maitra and Bhattacharya (2016c).

S-4 Proof of Lemma 11 of MB

For notational simplicity, let

$$\begin{aligned} \check{V}_{x^i, \beta_0}(s) &= \frac{b_{\beta_0}^2(X_i(s, x^i))}{\sigma^2(X_i(s, x^i))}; \\ \check{V}_{x^i, \beta_0, \beta_j}(s) &= \frac{b_{\beta_j}(X_i(s, x^i))b_{\beta_0}(X_i(s, x^i))}{\sigma^2(X_i(s, x^i))}; \\ \check{V}_{x^i, \beta_1}(s) &= \frac{b_{\beta_1}^2(X_i(s, x^i))}{\sigma^2(X_i(s, x^i))}. \end{aligned}$$

Continuity of $\mathcal{K}(f_{\theta_0, x, T, z}, f_{\theta_1, x, T, z})$ with respect to x and T , the fact that $x^k \rightarrow x^\infty$ and $T_k \rightarrow T^\infty$

as $k \rightarrow \infty$, assumption (H5'), and the dominated convergence theorem together ensure that

$$\begin{aligned}
& \lim_{n \rightarrow \infty} \frac{\sum_{k=1}^n \mathcal{K}(f_{\theta_0, x^k, T_k, z_k}, f_{\theta_1, x^k, T_k, z_k})}{n} \\
&= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \int_0^{T_k} \left\{ \frac{\phi_{k, \xi_0}(s)^2}{2} E_{\theta_0}(\check{V}_{x^k, \beta_0}(s)) \right. \\
&\quad \left. - \phi_{k, \xi_0}(s) \phi_{k, \xi_1}(s) E_{\theta_0}(\check{V}_{x^k, \beta_0, \beta_1}(s)) + \frac{\phi_{k, \xi_1}(s)^2}{2} E_{\theta_0}(\check{V}_{x^k, \beta_1}(s)) \right\} ds \\
&= \lim_{n \rightarrow \infty} \frac{1}{n} \int_0^{T_k} \sum_{k=1}^n \left\{ \frac{(\xi_{00} + \xi_{10}g_1(z_{k1}(s)) + \cdots + \xi_{p0}g_p(z_{kp}(s)))^2}{2} E_{\theta_0}(\check{V}_{x^k, \beta_0}(s)) \right. \\
&\quad - (\xi_{00} + \xi_{10}g_1(z_{k1}(s)) + \cdots + \xi_{p0}g_p(z_{kp}(s)))(\xi_{01} + \xi_{11}g_1(z_{k1}(s)) + \cdots + \xi_{p1}g_p(z_{kp}(s))) E_{\theta_0}(\check{V}_{x^k, \beta_0, \beta_1}(s)) \\
&\quad \left. + \frac{(\xi_{01} + \xi_{11}g_1(z_{k1}(s)) + \cdots + \xi_{p1}g_p(z_{kp}(s)))^2}{2} E_{\theta_0}(\check{V}_{x^k, \beta_1}(s)) \right\} ds \\
&= \int_0^{T^\infty} \left\{ \left(\frac{\xi_{00}^2}{2} + \xi_{00} \sum_{l=1}^p \xi_{l0} c_l(s) + \frac{1}{2} \sum_{l=1}^p \sum_{m=1}^p \xi_{l0} \xi_{m0} c_l(s) c_m(s) \right) E_{\theta_0}(\check{V}_{x^\infty, \beta_0}(s)) \right. \\
&\quad - \left(\xi_{00} \xi_{01} + \xi_{00} \sum_{l=1}^p \xi_{l1} c_l(s) + \xi_{01} \sum_{l=1}^p \xi_{l0} c_l(s) + \sum_{l=1}^p \sum_{m=1}^p \xi_{l0} \xi_{m1} c_l(s) c_m(s) \right) E_{\theta_0}(\check{V}_{x^\infty, \beta_0, \beta_1}(s)) \\
&\quad \left. + \left(\frac{\xi_{01}^2}{2} + \xi_{01} \sum_{l=1}^p \xi_{l1} c_l(s) + \frac{1}{2} \sum_{l=1}^p \sum_{m=1}^p \xi_{l1} \xi_{m1} c_l(s) c_m(s) \right) E_{\theta_0}(\check{V}_{x^\infty, \beta_1}(s)) \right\} ds, \tag{S-4.1}
\end{aligned}$$

which is the Kullback-Leibler distance between the models of the same form as $f_{\theta_0, x, T, z}$ and $f_{\theta_1, x, T, z}$, but with x, T and $g_l(z_l(s))$ replaced with x^∞, T^∞ and $c_l(s)$.

References

- Akaike, H. (1973). Information theory and an extension of the maximum likelihood principle. In B. N. Petrov and F. Csaki, editors, *Second International Symposium on Information Theory*, pages 267–281, Budapest. Akademiai Kiado. Reprinted in S. Kotz and N. L. Johnson (Eds) (1992). *Breakthroughs in Statistics Volume I: Foundations and Basic Theory*, pp. 610–624. Springer-Verlag.
- Barron, A., Schervish, M. J., and Wasserman, L. (1999). The Consistency of Posterior Distributions in Nonparametric Problems. *The Annals of Statistics*, **27**, 536–561.
- Berge, C. (1963). *Espaces Topologiques*. Dunod, Paris. Translated version: Topological Spaces. Translated by E. M. Patterson, Oliver and Boyd, Edinburgh.
- Delattre, M., Genon-Catalot, V., and Samson, A. (2013). Maximum Likelihood Estimation for Stochastic Differential Equations with Random Effects. *Scandinavian Journal of Statistics*, **40**, 322–343.
- Fuchs, C. (2013). *Inference for Diffusion Processes: With Applications in Life Sciences*. Springer, New York.
- Iacus, S. M. (2008). *Simulation and Inference for Stochastic Differential Equations: With R Examples*. Springer, New York.
- Jeffreys, H. (1961). *Theory of Probability*. 3rd edition. Oxford University Press, Oxford.
- Kass, R. E. and Raftery, R. E. (1995). Bayes factors. *Journal of the American Statistical Association*, **90**(430), 773–795.

- Leander, J., Almquist, J., Ahlström, C., Gabrielsson, J., and Jirstrand, M. (2015). Mixed Effects Modeling Using Stochastic Differential Equations: Illustrated by Pharmacokinetic Data of Nicotinic Acid in Obese Zucker Rats. *The AAPS Journal*, **17**, 586–596.
- Liu, J. (2001). *Monte Carlo Strategies in Scientific Computing*. Springer-Verlag, New York.
- Maitra, T. and Bhattacharya, S. (2015). On Bayesian Asymptotics in Stochastic Differential Equations with Random Effects. *Statistics and Probability Letters*, **103**, 148–159. Also available at “<http://arxiv.org/abs/1407.3971>”.
- Maitra, T. and Bhattacharya, S. (2016a). Asymptotic Theory of Bayes Factor in Stochastic Differential Equations. Submitted.
- Maitra, T. and Bhattacharya, S. (2016b). Asymptotic Theory of Bayes Factor in Stochastic Differential Equations: Part II. Submitted.
- Maitra, T. and Bhattacharya, S. (2016c). On Asymptotics Related to Classical Inference in Stochastic Differential Equations with Random Effects. *Statistics and Probability Letters*, **110**, 278–288. Also available at “<http://arxiv.org/abs/1407.3968>”.
- Maitra, T. and Bhattacharya, S. (2016d). Supplement to “Asymptotic Theory of Bayes Factor in Stochastic Differential Equations: Part I”. Submitted.
- Mao, X. (2011). *Stochastic Differential Equations and Applications*. Woodhead Publishing India Private Limited, New Delhi, India.
- Oravecz, Z., Tuerlinckx, F., and Vandekerckhove, J. (2011). A hierarchical latent stochastic differential equation model for affective dynamics. *Psychological Methods*, **16**, 468–490.
- Overgaard, R. V., Jonsson, N., Tornøe, C. W., and Madsen, H. (2005). Non-Linear Mixed-Effects Models with Stochastic Differential Equations: Implementation of an Estimation Algorithm. *Journal of Pharmacokinetics and Pharmacodynamics*, **32**, 85–107.
- Robert, C. P. and Casella, G. (2004). *Monte Carlo Statistical Methods*. Springer-Verlag, New York.
- Roberts, G. and Stramer, O. (2001). On Inference for Partially Observed Nonlinear Diffusion Models Using the Metropolis-hastings Algorithm. *Biometrika*, **88**, 603–621.
- Schwarz, G. (1978). Estimating the Dimension of a Model. *The Annals of Statistics*, **6**, 461–464.
- Sivaganesan, S. and Lingham, R. T. (2002). On the Asymptotic of the Intrinsic and Fractional Bayes Factors for Testing Some Diffusion Models. *Annals of the Institute of Statistical Mathematics*, **54**, 500–516.
- Walker, S. G. (2004). Modern Bayesian Asymptotics. *Statistical Science*, **19**, 111–117.
- Walker, S. G., Damien, P., and Lenk, P. (2004). On Priors With a Kullback-Leibler Property. *Journal of the American Statistical Association*, **99**, 404–408.